

NUMERICAL SIMULATION OF THE FRACTIONAL LANGEVIN EQUATION

by

Peng GUO, Changpin LI*, and Fanhai ZENG

Department of Mathematics, Shanghai University, Shanghai, China

Original scientific paper
DOI: 10.2298/TSCI110407073G

In this paper, we study the fractional Langevin equation, whose derivative is in Caputo sense. By using the derived numerical algorithm, we obtain the displacement and the mean square displacement which describe the dynamic behaviors of the fractional Langevin equation.

Key words: fractional Langevin equation, Riemann-Liouville derivative, Caputo derivative, mean square displacement, numerical simulation

Introduction

The fractional calculus has been studied for more than three hundred years. For a long time the fractional calculus has been studied only in the pure mathematical field. In history, Euler, Riemann, Liouville, Gruwald, Letnikov, Leibnitz, L'Hospital, *et al.*, contributed to the fractional calculus [1-3]. It has not attracted more attention because of not finding applications. However, in recent few decades, the fractional calculus has been widely used in many fields such as chaotic dynamics, viscoelasticity, acoustics, physical chemistry, electromagnetics, signal processing, earthquake prediction, *etc.* The study of fractional differential equations shifts from pure theory to real applications [2], [4-15]. Especially, the stochastic fractional differential equations have attracted increasing interests due to the potential applications. In effect, human evolutions are deterministic in local situations and in short terms but are random globally and in a long run. The stochastic fractional differential equation can reflect dual characters-stochasticity and globality. Hence, the stochastic fractional differential equation is possibly the better choice for characterizing human evolutions.

In this paper we will study the fractional Langevin equation where the fractional derivative is in Caputo sense. In 1908 the French physicist Langevin introduced the concept of the equation of motion with a random variable, which reads as:

$$m \frac{d^2 x(t)}{dt^2} = -\gamma \frac{dx(t)}{dt} + F(x) + \xi(t) \quad (1)$$

where m is the mass of the particle, γ – the coefficient of viscosity, $F(x)$ – the external force, and $\xi(t)$ – the random force. The Langevin equation is always regarded as the first stochastic differ-

* Corresponding author; e-mail: lcp@shu.edu.cn

ential equation. Although the classical Langevin equation has a fundamental role in so many areas such as physics, chemistry, signal processing, financial market, *etc.* There are still some dynamics, for example the anomalous diffusion (sub-diffusion and super-diffusion), power-law phenomena, long-tail character, long-range interaction, *etc.*, which can not be described by the classical Langevin equation. Therefore, the generalized Langevin equations have been introduced to model the above behaviors [16-23].

Among the generalized Langevin equations, the fractional version is often used, which is in the following form:

$$\frac{d^2 x(t)}{dt^2} = -\gamma \int_0^t (t-\tau)^{-\alpha} x'(\tau) d\tau + F(x) + \xi(t) \quad (2)$$

where $0 < \alpha < 1$, γ is a constant, $F(x)$ – an external force field, $\xi(t)$ – a random force, and $\langle \xi(t) \rangle = 0$, $\langle \xi(t) \xi(s) \rangle = \delta(t-s)$. Here, $\int_0^t (t-\tau)^{-\alpha} x'(\tau) d\tau = \Gamma(1-\alpha) {}_c D_{0,t}^{\alpha} x(t)$; the fractional derivative ${}_c D_{0,t}^{\alpha} x(t)$ will be defined in the following section.

Preliminaries

First of all, we give some basic definitions. In general, fractional calculus includes both fractional integral (integration) and fractional differentiation. The fractional integral mainly means the Riemann-Liouville integral. For fractional differentiation, however there exist more than six kinds of fractional derivatives. Among them, the Riemann-Liouville derivative and the Caputo derivative are mostly utilized. In the following, we only introduce the fractional integral, the Riemann-Liouville derivative and the Caputo derivative [2, 3, 8-10].

Definition 1. Fractional integral of function $f(x)$ with order $\alpha > 0$ is defined by:

$$D_{a,x}^{-\alpha} f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-\tau)^{\alpha-1} f(\tau) d\tau \quad (3)$$

Definition 2. Riemann-Liouville fractional derivative of function $f(x)$ with order $\alpha > 0$ is defined by:

$${}_R D_{a,t}^{\alpha} f(x) = \frac{1}{\Gamma(m-\alpha)} \frac{d^m}{dx^m} \int_a^x (x-\tau)^{m-\alpha-1} f(\tau) d\tau, \quad m-1 < \alpha < m \quad (4)$$

Definition 3. Caputo fractional derivative of function $f(x)$ with order $\alpha > 0$ is defined by:

$${}_c D_{a,t}^{\alpha} f(x) = \frac{1}{\Gamma(m-\alpha)} \int_a^x (x-\tau)^{m-\alpha-1} f^{(m)}(\tau) d\tau, \quad m-1 < \alpha < m \quad (5)$$

The above two fractional derivatives are not equivalent, whose important properties are listed below.

Property 2.1. Assume $m-1 < \alpha < m \in \mathbb{Z}^+$, $\beta > 0$, and that suitable smooth conditions are satisfied, $n \in \mathbb{Z}^+$. Then the following equalities hold:

$$\begin{aligned} {}_c D_{a,x}^{\alpha} f(x) &= {}_R D_{a,x}^{\alpha} \left[f(x) \sum_{k=0}^{m-1} \frac{f^{(k)}(a)}{\Gamma(k+1)} (x-a)^k \right] \\ {}_D_{a,x}^{-\alpha} {}_R D_{a,x}^{\alpha} f(x) &= f(x) - \sum_{k=0}^{m-1} \frac{[{}_R D_{a,x}^{\alpha-k} f(x)] a}{\Gamma(\alpha-k+1)} (x-a)^k \\ {}_R D_{a,x}^{\alpha} {}_R D_{a,x}^{-\beta} f(x) &= {}_R D_{a,x}^{\alpha-\beta} f(x) \end{aligned}$$

$${}_C D_{a,x}^\alpha f^{(n)}(x) = {}_C D_{a,x}^{\alpha+n} f(x)$$

where

$$[{}_{RL} D_{a,x}^{\alpha-k} f(x)](a) = [{}_{RL} D_{a,x}^{\alpha-k} f(x)]_{x=a}$$

Algorithm for the fractional Langevin equation

In this paper we will give the numerical simulation for the fractional Langevin equation (2).

Without external force

In the following, we consider the force free type, i. e. $F(x) = 0$. The above equation can be written as:

$$\frac{d^2 x(t)}{dt^2} = -\gamma \int_0^t (t-\tau)^{-\alpha} x'(\tau) d\tau + \xi(t) \quad (6)$$

then integrating the both sides of the above equation yields:

$$\begin{aligned} x'(t_{k+1}) - x'(t_k) = & -\frac{\gamma}{\Gamma(1-\alpha)} \int_0^{t_{k+1}} (t_{k+1}-\tau)^{-\alpha} x(\tau) d\tau + \frac{\gamma}{\Gamma(1-\alpha)} \int_0^{t_k} (t_k-\tau)^{-\alpha} x(\tau) d\tau + \\ & + \frac{\gamma x(0)}{\Gamma(1-\alpha)} (t_{k+1}^{1-\alpha} - t_k^{1-\alpha}) + \int_{t_k}^{t_{k+1}} dW(t) \end{aligned} \quad (7)$$

The first integral term and the second one in eq. (7) are approximated by the rectangle formula. For the first integral part in the r. h. s.:

$$\int_0^{t_{k+1}} (t_{k+1}-\tau)^{-\alpha} x(\tau) d\tau \approx \sum_{j=0}^k b_{j,k+1} x(t_j) \quad (8)$$

then we can calculate the coefficients:

$$b_{j,k+1} = \frac{h^{1-\alpha}}{1-\alpha} [(k+1-j)^{1-\alpha} - (k-j)^{1-\alpha}] \quad (9)$$

Similarly, for the second term in the r. h. s.:

$$\int_0^{t_k} (t_k-\tau)^{-\alpha} x(\tau) d\tau \approx \sum_{j=0}^{k-1} c_{j,k} x(t_j) \quad (10)$$

we can also get the coefficients:

$$c_{j,k} = \frac{h^{1-\alpha}}{1-\alpha} [(k-j)^{1-\alpha} - (k-j-1)^{1-\alpha}] \quad (11)$$

For the l. h. s. of eq. (7), we use the finite difference method to approximate it. Therefore we get the discrete formula:

$$\begin{aligned} x_{k+1} = & 2x_k - x_{k-1} - \frac{\gamma h}{\Gamma(1-\alpha)} \sum_{j=0}^k b_{k-j} y_j + \frac{\gamma h}{\Gamma(1-\alpha)} \sum_{j=0}^{k-1} b_{k-j-1} y_j - \\ & - \frac{\gamma h x(0)}{\Gamma(1-\alpha)} (t_k^{1-\alpha} - t_{k+1}^{1-\alpha}) + h[W(t_{k+1}) - W(t_k)], \quad k \geq 2 \end{aligned} \quad (12)$$

where $W(t)$ is a Wiener process with mean zero, and $dW(t)$ is in Ito sense, *i. e.*, an independent increment random. With the above algorithm we simulate its dynamic behaviors. The following figures give the displacement and the mean square displacement under different parameters. Here we take $x(0) = 0$, $x(1) = 0.001$.

Figures 1(a), 2(a), 3(a), and 4(a) indicate the displacement. Figures 1(b), 2(b), 3(b), and 4b indicate the mean square displacement. From the figures we find that the displacement depends heavily on the damping coefficient. If the damping coefficient is relatively smaller, the displacement is bigger; if the damping coefficient is relatively bigger, the displacement is smaller.




Figure 1. The numerical simulation of eq. (2), where $\alpha = 0.9$, $\gamma = 0.1$; (a) The displacement, (b) The mean square displacement




Figure 2. The numerical simulation of eq. (2), where $\alpha = 0.9$, $\gamma = 0.6$. (a) The displacement, (b) The mean square displacement




Figure 3. The numerical simulation of eq. (2), where $\alpha = 0.8, \gamma = 0.7$; (a) The displacement, (b) The mean square displacement




Figure 4. The numerical simulation of eq. (2), where $\alpha = 0.8, \gamma = 0.95$; (a) The displacement, (b) The mean square displacement

With constant external force

In the following part, we consider the fractional Langevin equation with a constant external force, i. e. $F(x) = F \neq 0$.

$$\frac{d^2x(t)}{dt^2} = -\gamma \int_0^t (t-\tau)^{-\alpha} x'(\tau) d\tau + F + \xi(t) \quad (13)$$

The algorithm is almost similar to that for the fractional Langevin equation without external force:

$$x_{k+1} = 2x_k - x_{k-1} - \frac{\gamma h}{\Gamma(1-\alpha)} \sum_{j=0}^k b_{k-j} x_j + \frac{\gamma h}{\Gamma(1-\alpha)} \sum_{j=0}^{k-1} b_{k-j-1} x_j - \frac{\gamma h x(0)}{\Gamma(1-\alpha)} (t_k^{1-\alpha} - t_{k+1}^{1-\alpha}) + h[W(t_{k+1}) - W(t_k)] + F_k h^2, \quad k \geq 2 \quad (14)$$

where

$$b_{j,k+1} = \frac{h^{1-\alpha}}{1-\alpha} [(k+1-j)^{1-\alpha} - (k-j)^{1-\alpha}] \quad (15)$$




Figure 5. The numerical simulation of eq. (2), where $F=2$, $\alpha=0.9$, $\gamma=0.6$; (a) The displacement, (b) The mean square displacement




Figure 6. The numerical simulation of equation (2), where $F=3$, $\alpha=0.8$, $\gamma=0.95$; (a) The displacement, (b) The mean square displacement

Then we display the figs with different parameters (figs. 5 and 6).

Conclusions

In this paper we study the generalized Langevin equation with a memory kernel in the damping item, *i. e.* the fractional Langevin equation in Caputo sense. We study two cases of the fractional Langevin equation (*i. e.*, without force and with constant external force). Finally we give an algorithm and numerical experiments with different parameters. From the numerical simulations, we find that the displacement is bigger if the coefficient of the damping item is relatively smaller and *vice versa*.

Acknowledgment

This work was partially supported by the National Nature Science foundation of China (10872119) and the Shanghai Leading Academic Discipline Project (S30104).

References

- [1] Oldham, K. B., Spanier, J., *The Fractional Calculus*, Academic Press, New York, USA, 1974
- [2] Podlubny, I., *Fractional Differential Equations*, Academic Press, New York, USA, 1999
- [3] Kilbas, A., Srivastava, H., Trujillo, J., *Theory and Applications of Fractional Differential Equations*, Elsevier Science Ltd., Amsterdam, The Netherlands, 2006
- [4] Das, S., *Functional Fractional Calculus for System Identification and Controls*, Springer-Verlag, Berlin, 2008
- [5] Gorenflo, R., *Fractional Calculus*, Springer-Verlag, New York, USA, 1997
- [6] Gorenflo, R., Mainardi, F., Random Walk Models for Space-Fractional Diffusion Process, *Fractional Calculus and Applied Analysis*, 1 (1998), 2, pp. 167-191
- [7] Hilfer, R., *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, 2000
- [8] Li, C. P., Dao, X. H., Guo, P., Fractional Derivatives in Complex Planes, *Nonlinear Analysis: TMA*, 71 (2009), 5-6, pp. 1857-1869
- [9] Li, C. P., Deng, W. H., Remarks on Fractional Derivatives, *Appl. Math. Comput.*, 187 (2007), 2, pp. 777-784
- [10] Li, C. P., Qian, D. L., Chen, Y. Q., On Riemann-Liouville and Caputo Derivatives, *Discrete Dynamics in Nature and Society*, 2011 (2011), article 562494, pp. 1-15
- [11] Li, C. P., Zhao, Z. G., Chen, Y. Q., Numerical Approximation of Nonlinear Fractional Differential Equations with Subdiffusion and Superdiffusion, *Comput. Math. Appl.*, 62 (2011), 3, pp. 855-875
- [12] Miller, K. S., Ross, B., *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley and Sons – Interscience Publication, New York, USA, 1993
- [13] Zaslavsky, G. M., *Hamiltonian Chaos and Fractional Dynamics*, Oxford University Press, Oxford, UK, 2005
- [14] Mainardi, F., Gorenflo, R., On Mittag-Leffler-Type Functions in Fractional Evolution Process, *J. Comput. Appl. Math.*, 118 (2000), 1-2, pp. 283-299
- [15] Mainardi, F., The Fundamental Solutions for the Fractional Diffusion-Wave Equation, *Appl. Math. Lett.*, 9 (1996), 6, pp. 23-28
- [16] Fa, K. S., Fractional Langevin Equation and Riemann-Liouville Fractional Derivative, *Eur. Phys. J. E*, 24 (2007), 2, pp. 139-143
- [17] Eab, C. H., Lim, S. C., Fractional Generalized Langevin Equation Approach to Single-File Diffusion, *Physica A*, 389 (2010), 13, pp. 2510-2521
- [18] Lutz, E., Fractional Langevin Equation, *Phys. Rev. E*, 64 (2001), 5, pp. 051106-051110
- [19] Fa, K. S., Generalized Langevin Equation with Fractional Derivative and Long-Time Correlation Function, *Phys. Rev. E*, 73 (2006), 6, pp. 061104-061115
- [20] Kobolev, V., Romanov, E., Fractional Langevin Equation to Describe Anomalous Diffusion, *Progr. Theor. Phys. Suppl.* (2000), 139, pp. 470-476
- [21] West, B. J., Picozzi, S., Fractional Langevin Model of Memory in Financial Market, *Phys. Rev. E*, 65 (2002), 3, pp. 037106-037120
- [22] Wang, K., Lung, C., Long-Time Correlation Effects and Fractal Brownian Motion, *Phys. Lett. A*, 151 (1990), 3-4, pp. 119-121
- [23] Lim, S. C., Li, M., Teo, L. P., Langevin Equation with Two Fractional Orders, *Phys. Lett. A*, 372 (2008), 42, pp. 6309-6320