

THE INFLUENCE OF SOLAR ACTIVITIES ON OCCURRENCE OF THE FOREST FIRES IN SOUTH EUROPE

by

**Milan M. RADOVANOVIĆ^a, Tomislav M. PAVLOVIĆ^b, Gorica B. STANOJEVIĆ^a,
Miško M. MILANOVIĆ^c, Mila A. PAVLOVIĆ^c, and Aleksandar R. RADIVOJEVIĆ^d**

^a Geographical Institute "Jovan Cvijić", Serbian Academy of Sciences and Arts, Belgrade, Serbia

^b Department of Physics, Faculty of Natural Sciences and Mathematics,
University of Nis, Nis, Serbia

^c Faculty of Geography, University of Belgrade, Belgrade, Serbia

^d Department of Geography, Faculty of Sciences and Mathematics, University of Nis, Nis, Serbia

Original scientific paper

DOI: 10.2298/TSCI130930036R

In mid-July 2012, hundreds of fires spread over forest stands in the south of Europe. Considering available satellite and meteorological data we have tried to determine a method using analogy, whether these fires were preceded by the intensified solar activity. The justification of this approach lies in the belief that it is impossible by direct or indirect activity of man to set fire at the same time to the sites that are geographically separated. In this paper we have tried to test the hypothesis claiming that charged particles coming to us from the Sun may be responsible for the number of forest fires. Unlike some other situations when the potential explanation could be a sudden influx of the protons of different energy ranges, in our study, there are electrons as carriers of the potential explanation. Many previous studies results have shown that there is statistically significant relationship between the processes on the Sun and certain movements of air masses in the atmosphere our planet. The presented results in this paper of the correlation relationships between the mean hourly flows of electrons and the mean hourly air temperatures in Belgrade and Rome and the analysis of the synoptic situation as a function of the suggested model so far, indicate that the number of fires that were occurring more than a week, from July 15, 2012, in the south of Europe, were caused by electrons coming from the Sun.

Key words: *solar activity, electrons, forest fires, South Europe*

Introduction

On 15th July 2012 the public was informed that 212 fires were recorded in Serbia on that day. After examining the satellite image (fig. 1) it proved that at that time fires were registered in the wider area of the Balkan Peninsula, Sicily, northern Turkey, and Ukraine. The next few days the fires occurred in Bosnia and Herzegovina to such an extent that the authorities requested military aid from Croatia in order to prevent further spreading of the fire. After 2-3 days the public was informed that fires on the border between Spain and France were out of control and there were also casualties. The Canary Islands, Madeira Island, and about 5,000 ha of forests in the south of Portugal in the Algarve tourist region were also endangered [1].

* Corresponding author; e-mail: milan.geograf@gmail.com



Figure 1. Satellite image of the distribution of fire over the Balkan Peninsula and adjacent areas 07/15/2012 09:25 UTC. Retrieved July 16 2012 from <http://rapidfire.sci.gsfc.nasa.gov/cgi-bin/imager/gallery.cgi>

In Croatia, near the town of Rijeka, a fire-fighter was killed. On 18th July Greek authorities declared the state of emergency in five villages near the city of Patra, in the northern part of the Peloponnese peninsula on the rough terrain due to the fires that raged for days. The suburb of Athens was also threatened by the destructive power of fire. In the coming days information on a number of sites affected by fire in Macedonia and Montenegro was continuously arriving. On 24th July, 2012 in the evening it started raining which considerably facilitated the situation in most of the Balkan Peninsula. It is important to note that many of the locations burning (hot spots), of smaller area of 1.1 km² are not registered on satellite images because of the limited possibilities of measuring instruments. As in many other situations studied the cause of the initial phase of the flame in this case also remains unknown. Gorte, points out that: "Research information on causative factors and on the complex circumstances surrounding wildfire is limited" [2]. The value of wildfires as case studies for building predictive models is confined because the *a priori* situation (*e. g.* fuel loads and distribution) and burning conditions (*e. g.* wind and moisture levels, patterns, and variations) are often unknown.

Even if the possible role of pyromaniacs (as well as any other potential explanation regarding the direct or indirect role of man) is accepted for the occurrence of fire in specific locations, it remains unclear how there is a selective ignition of vegetation in remote places on the same day [3, 4]. In Serbia the drought dominated a few weeks earlier so the assumption that electrical discharges in the atmosphere may be an explanation, has remained unsubstantiated. It is unrealistic to expect that lightning can be attributed to biomass burning at such remote locations without being rainfall in any of them [5, 6]. It is founded that from 1990 to 1998 over 17000 naturally ignited wildfires were observed in Arizona and New Mexico on US federal land during the fire season of April through October. Lightning strikes associated with these fires accounted for less than 0.35 % of all recorded cloud-to-ground lightning strikes that occurred during the fire season during that time [7]. In addition, the role of the wind in the spread of the fire is not clear, especially if one takes into account the *skip* of certain vegetation complexes horizontally as well as vertically, *i. e.* altitude [8]. The studies conducted by Csiszar *et al.*, [9] bring additional concerns about the coupling of climate and fires. According to them, the most numerous fires detected by satellites in the period 2001-2002 took place from November to February in the belt of 7.5 to 22.5° N. The problem of collecting data using uniform methodology greatly burdens not only any serious statistical spatial-temporal analysis but also the work on any kind of projection models [10]. That is why the following statement does not come on surprise: "Thus, research on fire protection and control is considering challenging and predictive tools for fire protection and

control are often based substantially on the expert opinion and anecdotes rather than on the documented research evidence" [11].

When the region of the south Europe it has been noticed that with the increase in average annual temperatures it comes to an increase in the number of fires in the past few decades. Regarding the region of the Mediterranean it is particularly necessary to point on that a recent regional situation analysis published in the frame of the FAO Global Forest Fire Assessment 1990-2000 reveal that the average annual number of forest fires throughout the Mediterranean basin is close to 50 000, *i. e.* twice as many as during the 1970 [12]. According to Nikolov [13] it can be concluded that, in average, 58.8% from the total number of forest fires in the country of the Balkan Peninsula for the period 1988-2004 has human origin, 3.3% has natural origin and 37.9% has unknown origin. Given the geographic spread of fires in the mentioned period we have tried in the paper to test the hypothesis according to which the forest fires, for which there are no determined causes, are connected with the processes on the Sun, that is, charged particles [14]. According to these authors, charged particles are able in certain circumstances to penetrate the ground and burn plant mass which actually represents the initial phase of flame phenomenon. Under certain conditions it is primarily meant on the critical level of density of particles per

unit volume which diffuse to the lower layers of the atmosphere. Protons and electrons (with a very strong eruptions, nucleons, too), ejected from the coronal holes and/or energy sources towards the Earth are moving in the form of an electric jet. A simplified representation of the penetration of the solar wind (SW) through the atmosphere is given in fig. 2. Increased cloudiness is also a limiting factor because the water vapour in the atmosphere acts as absorbent of the charged particles [15]. In the polar areas due to the SW burst a process known as re-connection takes place. If the B_z component of the interplanetary magnetic field (IMF) has a negative sign it comes to merging with the geomagnetic field, *i. e.* re-connection in the areas above the North Pole. In contrast, it comes to re-connection over the Antarctic if B_z component has a positive sign [16]. Thus, the process is going on in the area where the Earth's magnetic field is the strongest. Otherwise, the IMF goes around the Earth. "We found that the events occurring during closed geomagnetic conditions do not show common peaks at all high latitude stations and tend to be coherent only among the Antarctic stations, while there is a lack of coherence between high latitude opposite the hemispheres. Conversely, during open geomagnetic conditions the pulsation events are characterized by discrete frequencies, the same at all stations, and are generally highly coherent between high and low latitudes and between opposite the hemispheres" [17].

In the equatorial regions the penetration occurs due to the orientation of the IMF towards the part of the geomagnetic field which is the weakest. Theoretical model according to which the charged particles penetrate the ground through geomagnetic anomaly and after scattering on the ground could cause burning of biomass, for the first time, to our knowledge, was presented by Gomes and Radovanović [18]. At the same time there is an impression that these processes are accompanied with certain patterns of disorders in the movement of air masses




Figure 2. Schematic survey of the way of SW penetration towards topographic surface [13]

[19]. Idea on the dominant influence of the processes on the Sun (including cosmic rays) on the movement of air masses in the atmosphere is not new [20-24]. According to Troshichev and Janzhura [25], the disturbance in the fluctuation of the SW causes the changes in the atmospheric electric field resulting in the alterations of the cloudiness of troposphere and the atmospheric radiation budget and dynamics. Georgieva *et al.* [26] have come to the conclusion that long-periodic correlation between the solar activity and the atmospheric circulation changes in the consecutive solar secular cycles and it depends on the north-south asymmetry. McKenzie, *et al.* [27] showed that certain quantitative connection between fires and drought periods do exist in eastern Washington, as well as quasi-periodical connection with the El-Nino/Southern oscillation (ENSO) (3-7 years periodicity) and the Pacific (inter) decadal oscillation (PDO) (20-30 years periodicity).

On the basis of the available data applying complex linear regression for the period 1891-2004, Milovanović and Radovanović [28] tested the connection between the solar activity and atmospheric circulation. The values for adjusted R2 were calculated from 0.572-0.825. According to Tinsley and Yu [29] "there has not been any decisive result which would discern how many of the monitored decade variations were formed because of the entry of the flux of particles, comparing to the total or spectral changes of radiation. However, there is not such ambiguity concerning the correlation of the atmospheric dynamics with particle flows on the weather scale day after day." The report by Baldwin and Dunkerton [30] shows that stratospheric mean-flow variations induce circulations that penetrate into the lower troposphere. To investigate these results more in detail, Boberg [31] have used the geopotential height (GPH) data on 16 pressure levels covering both hemispheres to see if the proposed correlation exists in the terrestrial stratosphere and troposphere. The results show high statistical connection between solar wind electric field (E) and geopotential height (E-GPH statistical connection), extending from the lower stratosphere down to the surface.

Causality of processes on the Sun and forest fires in South-East Europe on July 15th, 2012

Based on the theoretical considerations as well as the analysis of processes in the interplanetary space and in the atmosphere, we will attempt to test the hypothesis by which the forest fires are preceded by scattering of charged particles through the lower layers of the troposphere. According to data taken from NASA (fig. 3, left) a strong transmission of energy from the Sun occurred on July 12th 2012. In the immediate vicinity of the active region 1520 there were regions 1519 and 1521 (http://www.nasa.gov/mission_pages/sunearth/news/News071212-X1.4flare.html). On the same day, in the afternoon, in the interplanetary space, a sudden increase in the flow of charged particles occurred in all energy ranges (fig. 3, right).

Figures 4 and 5 confirm visually that there is a potential connection between the sudden influx of the energy from the Sun and the geomagnetic anomaly. Following the images that were taken from the same source in the coming hours it was observed that there was also a re-connection.

When it comes to re-connection it appears that at the beginning the stronger flow of energy to the Earth has mainly been directed towards the southern hemisphere. The first serious intrusion on the northern hemisphere over the polar region occurred in 2012 07 15 at 0023 UT when the measured energy flow of 93.1 GW was measured (Retrieved July 16, 2012 from <http://www.swpc.noaa.gov/pmap/Plots.html>). On the same day at 1536 UT, the flow rate of 384.3 GW was measured (taken from the same source). Following the idea according to which a sudden influx of charged particles has the possibility of penetration into the lower layers of the





Figure 3. An X1.4 class flare erupted from the center of the Sun, peaking on July 12, 2012 at 12:52 PM EDT. It erupted from Active Region 1520 which rotated into view on July 6. Retrieved July 16, 2012 from (http://www.nasa.gov/mission_pages/sunearth/news/News071212-X1.4flare.html) (left) and a sudden rise of electron (upper sketch) and proton (bottom sketch) flows in all energy ranges on July 12th, 2012 reached a peak late in the evening on July 14th, 2012. Retrieved July 17 2012 from (http://www.swpc.noaa.gov/ace/EPAM_7d.html) (right) (for color image see journal web site)



Figures 4 and 5. Ionospheric conditions on July 12th, 2012 over the geomagnetic anomaly of the western Atlantic and eastern Pacific. Left image is related to 15:59 and right to 16:59 UTC. Retrieved July 17, 2012 from (http://www.ngdc.noaa.gov/stp/drap/data/2012/07/12/SWX_DRAP20_C_SWPC_20120712165900_GLOBAL.png) (for color image see journal web site)

troposphere an analysis of the movement of air masses was carried out on the day when numerous fires occurred in Serbia (fig. 6). Yellow isolines in the figure show an average wind speed both horizontally and vertically.

In fig. 6 we can see clearly how the wind speeds reached the highest value northwest of Portugal and Spain to the Pyrenees, the Alps, the northern Adriatic Sea, and the north-western and northern Balkan Peninsula. The maximum values were reaching over 80 knt (~ 40 m/s) and it should not be disregarded that this has been a curved movement of air masses of the approximate west-east direction, but also from above to downward. Jet Stream over Europe on 250 hPa and Synoptic situation over Europe on July 15th, 2012, 00 UTC are given in fig. 7.




Figure 6. Wind Shear in the 150-300 mb layer mean minus 700-925 mb layer mean. Retrieved July 16, 2012 from <http://tropic.ssec.wisc.edu/real>




Figure 7. Jet stream over Europe on 250 hPa. Retrieved July 15, 2012 from <http://weather.uwyo.edu/cgi-bin/uamap> (left) and Synoptic situation over Europe on July 15th, 2012, 00 UTC Retrieved July 16, 2012 from (<http://meteonet.nl/aktueel/brackall.htm>) (right)

Based on fig. 7 (left) we can also see a similar direction of movement of air masses (jet stream) except that, unlike the previous image, in this we can see 2d representation which confirms that there has been a strong movement of air masses over the same geographical area. In the days that followed an increased number of fires coincided with these atmospheric disturbances. According to Gabis and Troshichev [32], fluctuations of baric field within periods of 5 ± 10 days are typical of the meridional and zonal transfer in the troposphere (500 mb-level).

Based on the synoptic map (fig. 7, right) we can see that there is a relatively high air pressure from 1014-1015 mb over almost the whole of the Mediterranean and the Balkan Peninsula (including the areas to the Black Sea), while up to 1020 mb was measured over the Iberian Peninsula. In contrast, left of the main direction of the jet stream shown in figs. 7 and 8, in central and north Europe, there is a relatively low air pressure – below 1007 mb. Analysis which would aim to connect the atmospheric pressure with the scattering of charged particles in the northern part of Africa and north Europe would imply much more complex approach, which was not the goal of this research.

According to Black [33] the results are in keeping with the view where the potential anomalies of vortex in the lower part of the stratosphere, associated with changes in strength of the stratospheric polar whirlpool (vortex) are causing zonal symmetric wind disturbances, spreading down toward the surface. The position of the opening of the magnetospheric door greatly depends on the moving of the geomagnetic poles for it is clear the magnetosphere co-ordinates (including also geomagnetic anomalies) link magnetic poles, and not the geographic ones. This is an essential factor which should represent the integral part of the forecasting models [4]. Radovanović [34] emphasizes the fundamental importance of the south magnetic pole moving (which is located in the northern hemisphere). The moving of the poles is directly connected with the movement of the entire network of geomagnetic lines, including the positions of anomalies. Therefore, if there is a hydrodynamic intake of air masses within these processes, then their influence on the meteorological conditions should be followed by the movement of magnetospheric co-ordinate network. In order to come to more real conclusions it is clear that the temperature data must be examined in much greater database and the links with some other indicators of the solar activity. Nikolić *et al.*, [35] give arguments that connect the daily mean values of proton flux with the average daily values of air temperatures in Torino (Italy). A drop in temperature, at the field of low atmospheric pressure, is the consequence of the downward

vertical advection of cold air masses under the effect of the dynamic pressure of proton particles of corpuscular solar radiation, that is, shock wave protons. In contrast, areas that come under the influence of electrons, on the basis of presented considerations, should be characterized by relatively stable weather conditions and increased air pressure [36].

Based on this approach, in general, it is possible on the basis of detection of a sudden influx of charged particles to expect the occurrence of forest fires for up to several days in advance. However, for precise determination of location where the fires will occur, the research of experts from various fields is necessary. Such a team research should provide the answer to the question of how it comes to a propagation of charged particles to the ground, *i. e.* burning of bio mass.

The theoretical and mathematical considerations of the charged particles moving through the troposphere

Let us suppose that with a deeper penetration of the current field to the ground, the power of the electric field becomes weaker and thus the power of its magnetic layer, too. After opening, the charged particles begin to scatter from the current field on the principle of the left coil: protons to the left and electrons to the right in relation to the dominant direction of current field [37]. In fact the intensity and the direction of the electromagnetic force \vec{F} is determined by the vector product:

$$d\vec{F} = I d\vec{l} \times \vec{B} \quad (1)$$

where I is the electric convection current generated by particles in motion, $d\vec{l}$ – the vector of length of the current element, and \vec{B} – the magnetic induction strength vector.

Upper relation connects electrical and mechanical values with the magnetic ones. When we apply this relation to the free electric loads of the SW, ranging in geomagnetic field, then we can say that the electromagnetic force is essentially a physical force, acting on the free electrical loads:

$$\vec{F} = q\vec{v} \times \vec{B} \quad (2)$$

where q is the electrical load of particles, and \vec{v} – the particle velocity.

When a SW charged particle, which has a velocity v , electrical load q , and mass m , penetrates into the geomagnetic field of the induction B after opening the current field, then it is affected by the electromagnetic force, that is, the force of electric and magnetic fields.

Bearing in mind that the particle velocity \vec{v} covers an angle θ with the vector of the magnetic induction \vec{B} , the velocity can be decomposed into two components: component $v \cos \theta$, which is in the direction of the field and component $v \sin \theta$, which is perpendicular to the direction of the magnetic field. The first longitudinal component of the particle velocity $v \cos \theta$ indicates that the movement of particle will be even and in the direction of the magnetic field. Another transversal component of the particle velocity $v \sin \theta$ will cause circular movement in the level which is perpendicular to the magnetic field (fig. 8). By mutual action of these components one gets that the resultant particle trajectory is coil with a cylinder-shaped tube, with a radius

$$r = \frac{mv \sin \theta}{qB} \quad (3)$$

$$d = \left(\frac{2\pi r}{v \sin \theta} \right) v \cos \theta = \frac{2\pi r m v \cos \theta}{qB} \quad (4)$$




Figure 8. Schematic representation of the resultant trajectory of the corpuscular radiation particle

This means that with deeper penetration into the lower layers of the atmosphere, the influence of geomagnetic field increases, that is, geomagnetic induction B increases, which results in a decrease in the radius of the current field. Magnetic layer of the current field does not allow the scattering of particles and with decreasing radius particle density increases, that is, momentum.

Wind speed outside the equatorial belt can be described by

$$v = c \frac{rqB}{m \sin \theta} \quad (5)$$

where c is the sliding factor

Momentum \vec{p} is a vector value, which intensity is defined by product of vector \vec{v} and scalar m , where \vec{v} – the velocity, and m – the mass of particles.

$$\vec{p} = m\vec{v} \quad (6)$$




Figure 9. Schematic representation of the downward vertical advection

The speed of change of momentum in time is equal to the force acting and has the same direction as the force (fig. 9).

Distribution of particles through the lower layers of the troposphere will depend on the altitude and the angle at which scattering of particles occurs. At the moment of loss of circulation kinetic energy the gravity and electric field force will act upon electrons. If the loss of kinetic energy of electrons occurs at relatively higher altitudes, the impact of electric field force will prevail. Otherwise, the electrons will be directed towards the ground.

Analysis of the flux of protons and electrons with series of air temperature and humidity in Belgrade and Rome in the period July 9th-24th, 2012

Considering the previous results we have tried to establish a quantitative relationship between hourly flow rates of electrons and protons and hourly values of air temperature and relative humidity in Belgrade and Rome for a period July 09-24, 2012 (Data for hourly flow rates of protons and electrons are taken from: http://www.swpc.noaa.gov/ftplib/lists/ace2/201207_ace_epam_1h.txt. Data for daily flow rates of protons and electrons are taken from: <http://www.swpc.noaa.gov/weekly/index.html>. Data for hourly values of air temperature and relative humidity in Belgrade were obtained from the Republic Hydrometeorological Service of Serbia. The daily average temperatures in Rome were kindly send by AIR FORCE, C.N.M.C.A., (National Centre of Meteorology, Air and Clima, 3° Service "Climatology", Rome, Italy). We analyzed the period from the beginning of the rapid flow of charged particles to July 24th when the rain subsided the situation with the fires in the Balkan Peninsula. Data for the station of Belgrade were measured according to the local time hence it was necessary to be complied with the co-ordinated universal time (UTC). In the analysis, 2-hour time lag of time series in Belgrade was taken into account, compared to the measurements of protons and electrons. We assumed

that it takes a certain period of time while charged particles reach the lower elevations and eventually are reflected on the down-to-earth measurements. Looking strictly mathematically it is found that the calculated values are relatively low: the strongest correlation was obtained for the flow of electrons in the range of 38-53 particles/cm²-s-ster-MeV and an air temperature of $r = 0.29$. In view of a possible link between the flow of charged particles and relative air humidity the strongest relationship was obtained for the flux of electrons in the same range but the negative correlation of $r = -0.32$ (tab. 1).

Table 1. Pearson correlation coefficients for hourly and daily values of electrons and protons and temperature (T) and relative humidity (R) in Belgrade, period July 9-24, 2012

Hourly values	T	R	Daily values	T	R
Electrons 38-53/cm ² -s-ster-MeV	0.29*	-0.32*	Electrons 38-53/cm ² -s-ster-MeV	0.34	-0.16
Electrons 175-315/cm ² -s-ster-MeV	0.10**	-0.17*	Electrons 175-315/cm ² -s-ster-MeV	0.14	0.17
Protons 47-68/cm ² -s-ster-MeV	0.20*	-0.22*	Protons 47-68/cm ² -s-ster-MeV	0.19	-0.20
Protons 115-195/cm ² -s-ster-MeV	0.23*	-0.23*	Protons 115-195/cm ² -s-ster-MeV	0.22	-0.20
Protons 310-580/cm ² -s-ster-MeV	0.25*	-0.25*	Protons 310-580/cm ² -s-ster-MeV	0.24	-0.19
Protons 795-1193/cm ² -s-ster-MeV	0.25*	-0.26*	Protons 795-1193/cm ² -s-ster-MeV	0.23	-0.14
Protons 1060-1900/cm ² -s-ster-MeV	0.26*	-0.26*	Protons 1060-1900/cm ² -s-ster-MeV	0.24	-0.13

* Significant at 0.05 level; ** Significant at 0.10 level

At the start of the study it was clear that strong correlations cannot be expected, primarily due to the large range of charged particles flow in a relatively short time compared to the values of air temperature and relative humidity. Despite the fact that this is a statistically small sample, testing is done on correlation links at daily averaged values. The obtained results are slightly better for the flux of electrons in the range of 38-53 particles/cm²-s-ster-MeV and air temperature $r = 0.34$ (with data for the station of Belgrade (tab. 1) and the flux of protons $p > 100$ MeV and relative air humidity $r = 0.56$ (tab. 2). In the analysis for Rome (weather station AM Rome Ciampino Airport), we had daily mean values of air temperature (tab. 3). It turned out that almost identical values were obtained for the flux of electrons in the range of 38-53 particles/cm²-s-ster-MeV and air temperature as for Belgrade $r = 0.37$, while for the flux of protons $p > 100$ MeV and the air temperature somewhat different values of $r = -0.58$ were obtained.

Table 2. Pearson correlation coefficients for daily values of electrons and protons (different scale for flux) and temperature (T) and relative humidity (R) in Belgrade, period 9-24 July 2012

	T	R
Electrons >2/cm ² -s-ster-MeV	0.17	-0.23
Protons >1/cm ² -s-ster-MeV	0.21	-0.21
Protons >10/cm ² -s-ster-MeV	0.21	-0.22
Protons >100/cm ² -s-ster-MeV	0.05	0.56*


* Significant at 0.05 level

For a more detailed consideration of the causal connection between the propagation of protons and electrons to the ground and the occurrence of forest fires it is necessary to bear in mind that penetration of the current field through the troposphere is not uniform. Therefore, the effect that charged particles would cause on the ground requires more complex modelling. In fig. 10 we can see that there are certain *matches* in the example of hourly mean values of air temperature in Belgrade and the flow of electrons in the range 175-315/cm²-s-ster-MeV.

Table 3. Pearson correlation coefficients for daily values of electrons and protons and temperature (T) in Rome, period July 9-24, 2012 and Pearson correlation coefficients for daily values of electrons and protons (different scale for flux) and temperature (T) in Rome, period July 9-24 2012

Correlation coefficients for daily values of electrons and protons and temperature (T) in Rome, period July 9-24, 2012		Correlation coefficients for daily values of electrons and protons (different scale for flux) and temperature (T) in Rome, period July 9-24, 2012	
	T		T
Electrons 38-53/cm ² -s-ster-MeV	0.37	Electrons >2/cm ² -s-ster-MeV	0.28
Electrons 175-315/cm ² -s-ster-MeV	-0.07	Protons >1/cm ² -s-ster-MeV	0.38
Protons 47-68/cm ² -s-ster-MeV	0.37	Protons >10/cm ² -s-ster-MeV	0.29
Protons 115-195/cm ² -s-ster-MeV	0.38	Protons >100/cm ² -s-ster-MeV	-0.58*
Protons 310-580/cm ² -s-ster-MeV	0.38	–	–
Protons 795-1193/cm ² -s-ster-MeV	0.35	–	–
Protons 1060-1900/cm ² -s-ster-MeV	0.35	–	–

* Significant at level 0.05

**Figure 10. Hourly values of air temperature in Belgrade and electrons 175-315/cm²-s-ster-MeV in the period July 9-24, 2012**

Conclusions

In this paper using the method of analogy, and on the basis of the available satellite and meteorological material we have tried to determine whether the increased solar activity was preceded by the numerous forest fires in South Europe in mid-July 2012. In the period from July 15th most countries around the Balkan Peninsula issued a statement about the fires in their territories the situation was only claimed with the absence of rain on July 24th. The starting hypothesis is that the charged particles from the Sun are in certain circumstances able to penetrate the ground and burn plant mass, that is, initiate a flame. A strong emission of energy from the Sun occurred on July 12th, which was manifested in a sharp increase in the charged particles in all energy ranges in interplanetary space, in order that a maximum followed on July 14th, in the evening. Analysis of synoptic conditions on July 15th indicates increased wind speed (over 40 m/s) in the area above the northern part of the Iberian Peninsula, the Alps, the northern Adriatic Sea and the north-western and northern part of the Balkan Peninsula (curved moving of approximate west-east direction, and from above to downward), which coincides with an increased number

According to Weng [38], it should be kept in mind that once a solar preferred atmospheric circulation patterns and related weather/climate events occur, the non-linear wave-mean interaction will be in effect, resulting in zonal index cycle, which may or may not synchronize with solar activity.

In this regard Ogurtsov *et al.*, [39] point out that it can be very difficult to reveal these variations because the character of the originating cause-effect relation is extremely complex, and this problem can hardly be solved using the traditional methods of statistical analysis.

of forest fires in the coming days. At the same time, across the Mediterranean and the Balkan Peninsula there is relatively high air pressure (1014-1015 mb), which is consistent with the hypothesis that these areas were under the dominant influence of the electrons (right side of the jet stream main direction), that is, low pressure over northern and central Europe indicates the propagation of protons in these areas (left of the jet stream main direction).

Analysis of the quantitative agreement (in the period July 09-24, 2012) between the proton and electron flow and hourly values of the temperature and relative humidity in Belgrade, that is, mean daily temperatures in Rome, from a strictly mathematical point of view show relatively poor values. However, strong correlations cannot be expected given the large range of the charged particles flow in a relatively short time compared to the values of air temperature and relative humidity. Also, penetration of the current field through the troposphere is not uniform, and the effect of the charged particles on the ground requires much complex modelling. The obtained results can be a starting point for future forecasting models which for the announcement of fire elementals must contain in it the propagation of protons and electrons through the lower layers of the atmosphere.

Acknowledgment

The results are a part of the project III47007 and OI 176008 funded by the Ministry of Education, Science and Technological Development of the Republic of Serbia.

References

- [1] Rowell, A., Moore, F. P., Global Review of Forest Fires, WWF, IUCN, 2000
- [2] Gorte, W. R., Forest Fire/Wildfire Protection, CRS Report for Congress (Received through the CRS Web), Congressional Research Service, The Library of Congress, 2006, Order Code RL30755
- [3] Ducić, V., et al., Contemporary Climate Variability and Forest Fires in Deliblatska Peščara. *J. Geogr. Inst. Cvijić*, 58 (2008), pp. 59-73
- [4] Radovanović, M., et al., Endangerment of Undeveloped Areas of Serbia by Forest Fires, *J. Geographical Institute "J. Cvijić"*, 59 (2009), 2, pp. 17-35
- [5] Kourtz, P. H., Todd, J. B., Predicting the Daily Occurrence of Lightning-Caused Forest Fires, Forestry Canada Information Report, Petawawa National Forestry Institute, 1991, PI-X-112
- [6] Krawchuk, A. M., et al., Biotic and Abiotic Regulation of Lightning Fire Initiation in the Mixed Wood Boreal Forest, *Ecology*, 87 (2006), 2, pp. 458-468
- [7] Hall, L. B., Precipitation Associated with Lightning-Ignited Wildfires in Arizona and New Mexico, *Int J Wildland Fire*, 16 (2007), 2, pp. 242-254
- [8] Linn, R. R., Cunningham, Ph., Numerical Simulations of Grass Fires Using a Coupled Atmosphere-Fire Model: Dynamics of Fire Spread, *J Geophys Res.*, 110 (2005), 13, pp. 1-14
- [9] Csizsar, I., et al., Global Fire Activity from Two Years of MODIS Data, *Int J Wildland Fire*, 14 (2005), 2, pp. 117-130
- [10] Conard, G. S., et al., Determining Effects of Area Burned and Fire Severity on Carbon Cycling and Emissions in Siberia, *Climatic Change*, 55 (2002), 1-2, pp. 197-211
- [11] Gorte, W. R., Forest Fire Protection, CRS Report for Congress (Received through the CRS Web), Congressional Research Service, The Library of Congress, 2000; Order Code RL30755
- [12] Goldammer, G. J., Towards International Cooperation in Managing Forest Fire Disasters in the Mediterranean Region, *International Forest Fire News*, No. 27, July-December 2002, pp. 81-89
- [13] Nikolov, N., Global Forest Resources Assessment 2005 – Report on Fires in the Balkan Region, Forestry Department, FAO of the UN, Fire Management Working Papers FM/11/E, 2006, Rome (www.fao.org/forestry/site/fire-alerts/en).
- [14] Stevančević, M., et al., Solar Wind and the Magnetospheric Door as Factor of Atmospheric Processes, *Proceedings, Second International Conference Global Changes and New Challenges of 21st Century*, Sofia, Bulgaria, 2006, pp. 88-94
- [15] Mukherjee, S., Radovanović, M., Influence of the Sun in the Genesis of Tornadoes, *The IUP Journal of Earth Sciences*, 5 (2011), 1, pp. 7-21

- [16] Radovanović, M., et al., A Contribution to the Study of the Influence of the Energy of Solar Wind upon the Atmospheric Processes, *J. Geogr. Inst. Cvijic*, 53 (2003), pp. 1-18
- [17] Lepidi, S., et al., The Earth's Passage of Coronal Mass Eject on October 29-31, 2003: ULF Geomagnetic Field Fluctuations at Very High Latitude, *Memorie della Società Astronomica Italiana*, 76 (2005), 4, pp. 998-1001
- [18] Gomes, J. F. P., Radovanović, M., Solar Activity as a Possible Cause of Large Forest Fires – A Case Study: Analysis of the Portuguese Forest Fires, *Sci Total Environ*, 394 (2008), 1, pp. 197-205
- [19] Pereira, M. G., et al., Synoptic Patterns Associated with Large Summer Forest Fires in Portugal, *Agr Forest Meteorol*, 129 (2005), 1-2, pp. 11-25
- [20] Komitov, B. P., The Sun, Climate and Their Changes in Time, *Nauka*, XV, (2005), 6, pp. 28-39
- [21] Liliensten, J., Bornarel, J., *Space Weather, Environment and Societies*, Springer Ltd. 2006, DOI:10.1007/1-4020-4332-5
- [22] Mukherjee, S., Cosmic Influence on Sun-Earth Environment, *Sensors*, 8 (2008), 12, pp. 7736-52
- [23] Radovanović, M., Solar Activity – Climate Change and Natural Disasters in Mountain Regions, in: *Sustainable Development in Mountain Regions* (Chapter 2), Springer Science+Business Media B.V. 2011, pp. 9-17
- [24] Verdon, C. D., et al., Multi-Decadal Variability of Forest Fire Risk – Eastern Australia, *Int J Wildland Fire*, 13 (2004), 2, pp. 165-171
- [25] Troshichev, O. A., Janzhura, A., Temperature Alterations on the Antarctic Ice Sheet Initiated by the Disturbed Solar Wind, *J Atmos Sos-Terr Phy*, 66 (2004), 13-14, pp. 1159-72
- [26] Giorgieva, K., et al., Long-Term Variations in the Correlation between NAO and Solar Activity: The Importance of North-South Solar Activity Asymmetry for Atmospheric Circulation, *Adv Space Res*, 40 (2007), 7, pp. 1152-66
- [27] McKenzie, D., et al., Fire and Climatic Variability in the Inland Pacific Northwest: Integrating Science and Management, Final report to the Joint Fire Science Program on Project #01-1-6-01, U. S. Dept., of Agriculture, Pacific Northwest Research Station, 2004
- [28] Milovanović, B., Radovanović, M., The Connections between Solar Activity and the Circulation of Atmosphere in the 1891-2004 period (in Serbian), *J. Geogr. Inst. Cvijic*, 59 (2009), 1, pp. 35-48
- [29] Tinsley, A. B., Yu, F., Atmospheric Ionization and Clouds as Links between Solar Activity and Climate, in: AGU Monograph: Solar Variability and Its Effects on the Earth's Atmospheric and Climate System, AGU press, Washington, DC, 2004, pp. 321-340
- [30] Baldwin, M. P., Dunkerton, T. J., Stratospheric Harbingers of Anomalous Weather Regimes, *Science*, 294 (2001), 5542, pp. 581-584
- [31] Boberg F., Solar Wind Electric Field Modulation of the NAO: A Correlation Analysis in the Lower Atmosphere, *Geophys Res Lett*, 30 (2003), 15, p. 1825
- [32] Gabis, I. P., Troshichev, O. A., Influence of Short-Term Changes in Solar Activity on Baric Field Perturbations in the Stratosphere and Troposphere, *J Atmos Sol-Terr Phy*, 62 (2000), 9, pp. 725-735
- [33] Black, R. X., Stratospheric Forcing of Surface Climate in the Arctic Oscillation, *J. Climate*, 15 (2002), 3, pp. 268-277
- [34] Radovanović, M., b) The Basic Settings of Heliocentric Climate Model, Book of Abstracts, International Conference Regional Responses to Global Environmental Change in North-East and Central Asia, Plenary Session, Irkutsk, Russia, 2012, (1); pp. 52-54
- [35] Nikolić, J., et al., An Astrophysical Analysis of Weather Based on the Solar Wind Parameters, *Nucl Technol Radiat*, 25 (2010), 3, pp. 171-178
- [36] Radovanović, M., et al., Heliocentric Electromagnetic Long-Term Weather Forecast and its Applicable Significance, *J. Geogr. Inst. Cvijic*, 54 (2005), 1, pp. 5-18
- [37] Radovanović, M., Forest Fires in Europe from July 22nd to 25th, 2009, *Arch. Biol. Sci.*, 62 (2010), 2, pp. 419-24
- [38] Weng, H., Impacts of Multi-Scale Solar Activity on Climate, Part I: Atmospheric Circulation Patterns and Climate Extremes, *Adv Atmos Sci.*, 29 (2012), 4, pp. 867-886
- [39] Ogurtsov, M. G., et al., Possible Manifestation of Nonlinear Effects when Solar Activity Affects Climate Changes, *Geomagnetism and Aeronomy*, 50 (2010), 1, pp. 15-20

Paper submitted: September 30, 2013

Paper revised: March 20, 2014

Paper accepted: March 27, 2014