

FRACTIONAL DERIVATIVES OF CONSTANT AND VARIABLE ORDERS APPLIED TO ANOMALOUS RELAXATION MODELS IN HEAT TRANSFER PROBLEMS

by

Xiao-Jun YANG^{a,b,*}

^a State Key Laboratory for Geo-Mechanics and Deep Underground Engineering,
School of Mechanics and Civil Engineering, China University of Mining and Technology,
Xuzhou, China

^b School of Mechanics and Civil Engineering, China University of Mining and Technology,
Xuzhou, China

Original scientific paper
<https://doi.org/10.2298/TSCI161216326Y>

In this paper, we address a class of the fractional derivatives of constant and variable orders for the first time. Fractional-order relaxation equations of constants and variable orders in the sense of Caputo type are modeled from mathematical view of point. The comparative results of the anomalous relaxation among the various fractional derivatives are also given. They are very efficient in description of the complex phenomenon arising in heat transfer.

Key words: heat transfer, fractional derivative of constant, anomalous relaxation, fractional derivative of variable order, fractional differential equation

Introduction

Fractional derivatives (FD) were utilized to model the complex phenomenon in science and engineering practice [1-7]. The FD of constant order, *e. g.*, Riemann-Liouville and Caputo FD, were considered to describe the anomalous relaxation [2, 5, 8-10]. The Riemann-Liouville type FD of variable-order was proposed in [11, 12] and its extended versions were considered in [13]. The Caputo type FD of variable-order was developed in [14] and its extended versions were reported in [15, 16].

Recently, FD involving the kernels of the exponential and Mittag-Leffler functions were reported in [17-28]. For example, FD with respect to the kernel of the exponential function in the sense of the Caputo type was firstly reported in [17-24] and FD with the aid of the kernel of the exponential function in the sense of the Riemann-Liouville type was proposed in [25] and developed in [26]. The FD with the help of the kernel of the stretched exponential function in the sense of the Caputo type was presented in [27]. The FD containing the Mittag-Leffler function kernels in the sense of the Caputo and Riemann-Liouville types were developed in [28].

Due to the previous FD in the different kernels, a class of the FD of constant and variable orders, such as has not been reported. Motivated by the previous ideas, the brief aims of the previous paper are to suggest FD of constant order with respect to the Mittag-Leffler function and stretched Mittag-Leffler and exponential functions kernels in the sense of Gaussian-like type, the variable-order FD in the Caputo-Fabrizio, Sun-Hao-Zhang-Baleanu, and Atanga-

* Author's e-mail: dyangxiaojun@163.com

na-Baleanu types as well as Mittag-Leffler and exponential functions and their stretched and generalized versions and to present their applications for handling the anomalous relaxation models in heat transfer problems.

The FD of constant and variable orders

Let $\Xi(\mu)$ be the differentiable function defined on the interval $[a, b]$, as well as $\mathfrak{Z}(\omega)$ and $\mathfrak{Z}[\omega(x)]$ be normalization constants with respect to the orders ω ($0 \leq \omega \leq 1$) and $\omega(x)$ [$0 \leq \omega(x) \leq 1$], respectively. For more details of the normalization constants, readers refer to [17-28].

The FD of constant order

The FD of the Caputo type with respect to the singular power-law kernel is given by [2, 5, 10]:

$$D_{a^+}^{(\omega)} \Xi(x) = \frac{1}{\Gamma(1-\omega)} \int_a^x \frac{1}{(x-\mu)^\omega} \left[\frac{d\Xi(\mu)}{d\mu} \right] d\mu \quad (1)$$

where $a \leq x$, n is an integer, and $\Gamma(\cdot)$ denotes the gamma function.

The FD of the Caputo-Fabrizio type with respect to the exponential kernel, denoted by [17-23]:

$$y = \varphi_1(x, \omega) = \exp\left(-\frac{\omega}{1-\omega}x\right) = \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{x^i}{\Gamma(1+i)} \quad (2a)$$

is given by:

$${}^{\text{CF}}D_x^{(\omega)} \Xi(x) = \frac{(2-\omega)\mathfrak{Z}(\omega)}{2(1-\omega)} \int_a^x \exp\left[-\frac{\omega}{1-\omega}(x-\mu)\right] \Xi^{(1)}(\mu) d\mu \quad (2b)$$

where $a \leq x$.

For $\mathfrak{Z}(\omega) = 2/(2-\omega)$, eq. (2b) is given by [23]:

$${}^{\text{LNCF}}D_x^{(\omega)} \Xi(x) = \frac{1}{1-\omega} \int_0^x \exp\left[-\frac{\omega}{1-\omega}(x-\mu)\right] \Xi^{(1)}(\mu) d\mu \quad (3)$$

The FD of the Sun-Hao-Zhang-Baleanu type with respect to the stretched exponential function, denoted by:

$$y = \varphi_2(x, \omega) = \exp\left(-\frac{\omega}{1-\omega}x^\omega\right) = \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{x^{i\omega}}{\Gamma(1+i)} \quad (4a)$$

is given by [27]:

$${}^{\text{SHZB}}D_x^{(\omega)} \Xi(x) = \frac{\mathfrak{Z}(\omega)}{(1-\omega)^{1/\omega}} \int_a^x \exp\left[-\frac{\omega}{1-\omega}(x-\mu)^\omega\right] \Xi^{(1)}(\mu) d\mu \quad (4b)$$

where $a \leq x$.

For simplicity, we have [27]:

$$\mathfrak{Z}(\omega) = \Gamma(\omega+1) \quad (5a)$$

such that:

$${}^{\text{SHZB}}D_x^{(\omega)}\Xi(x) = \frac{\Gamma(1+\omega)}{(1-\omega)^{1/\omega}} \int_0^x \exp\left[-\frac{\omega}{1-\omega}(x-\mu)^\omega\right] \Xi^{(1)}(\mu) d\mu \quad (5b)$$

The FD with respect to the Mittag-Leffler function kernel, denoted by:

$$y = \varphi_3(x, \omega) = E_\omega\left(-\frac{\omega}{1-\omega}x\right) = \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{x^i}{\Gamma(i\omega+1)} \quad (6a)$$

is defined by:

$${}^{\text{YC}}D_x^{(\omega)}\Xi(x) = \frac{\mathfrak{I}(\omega)}{1-\omega} \int_a^x E_\omega\left[-\frac{\omega}{1-\omega}(x-\mu)\right] \Xi^{(1)}(\mu) d\mu \quad (6b)$$

where $a \leq x$.

The FD of the Atangana-Baleanu type with respect to the stretched Mittag-Leffler function kernel, denoted by:

$$y = \varphi_4(x, \omega) = E_\omega\left(-\frac{\omega}{1-\omega}x^\omega\right) = \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{x^{i\omega}}{\Gamma(i\omega+1)} \quad (7a)$$

is given by [28]:

$${}^{\text{ABC}}D_x^{(\omega)}\Xi(x) = \frac{\mathfrak{I}(\omega)}{1-\omega} \int_a^x E_\omega\left[-\frac{\omega}{1-\omega}(x-\mu)^\omega\right] \Xi^{(1)}(\mu) d\mu \quad (7b)$$

where $a \leq x$.

The FD of the Caputo-Fabrizio type with respect to the Gaussian-function kernel, denoted by:

$$y = \varphi_5(x, \omega) = \exp\left(-\frac{\omega}{1-\omega}x^2\right) = \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{x^{2i}}{\Gamma(1+i)} \quad (8a)$$

is given by [24]:

$${}^{\text{CF}}D_x^{(\omega)}\Xi(x) = \frac{1+\omega^2}{\sqrt{\pi^\omega(1-\omega)}} \int_a^x \exp\left[-\frac{\omega}{1-\omega}(x-\mu)^2\right] \Xi^{(1)}(\mu) d\mu \quad (8b)$$

where $a \leq x$.

The FD with respect to the stretched exponential function kernel in the sense of Gaussian-like type, denoted by:

$$y = \varphi_6(x, \omega) = \exp\left(-\frac{\omega}{1-\omega}x^{2\omega}\right) = \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{x^{2\omega i}}{\Gamma(1+i)} \quad (9a)$$

is defined by:

$${}^{\text{GYC}}D_x^{(\omega)}\Xi(x) = \frac{1+\omega^2}{\sqrt{\pi^\omega(1-\omega)}} \int_a^x \exp\left[-\frac{\omega}{1-\omega}(x-\mu)^{2\omega}\right] \Xi^{(1)}(\mu) d\mu \quad (9b)$$

where $a \leq x$.

The FD with respect to the stretched Mittag-Leffler function kernel in the sense of Gaussian-like type, denoted by:

$$y = \varphi_7(x, \omega) = E_\omega \left(-\frac{\omega}{1-\omega} x^{2\omega} \right) = \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega} \right)^i \frac{x^{2\omega i}}{\Gamma(1+i\omega)} \quad (10a)$$

is defined by:

$${}^{\text{GYGC}}D_x^{(\omega)}\Xi(x) = \frac{1+\omega^2}{\sqrt{\pi^\omega(1-\omega)}} \int_a^x E_\omega \left[-\frac{\omega}{1-\omega} (x-\mu)^{2\omega} \right] \Xi^{(1)}(\mu) d\mu \quad (10b)$$

where $a \leq x$.

The LT of the functions $x^{-1-\omega}/\Gamma(-\omega)$ and $E_\omega(-x^\omega)$ are given in [2, 12]:

$$\hat{L} \left[\frac{x^{-1-\omega}}{\Gamma(-\omega)} \right] = s^\omega \quad (11)$$

$$\hat{L} \left[E_\omega(-\beta x^\omega) \right] = \frac{s^{\omega-1}}{s^\omega + \beta} \quad (12)$$

respectively, where \hat{L} is the LT operator with respect to x , and β is a constant.

The LT of FD are:

$$\hat{L} \left[{}^{\text{SHZB}}D_x^{(\omega)}\Xi(x) \right] = \frac{\Im(\omega)}{(1-\omega)^{1/\omega}} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega} \right)^i \frac{\Gamma(1+i\omega)}{\Gamma(1+i)} \frac{[s\hat{\Xi}(s) - \Xi(0)]}{s^{i\omega+1}} \quad (13)$$

$$\hat{L} \left[{}^{\text{YC}}D_x^{(\omega)}\Xi(x) \right] = \frac{\Im(\omega)}{1-\omega} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega} \right)^i \frac{\Gamma(i+1)}{\Gamma(i\omega+1)} \frac{[s\hat{\Xi}(s) - \Xi(0)]}{s^{i+1}} \quad (14)$$

$$\hat{L} \left[{}^{\text{GYC}}D_x^{(\omega)}\Xi(x) \right] = \frac{1+\omega^2}{\sqrt{\pi^\omega(1-\omega)}} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega} \right)^i \frac{\Gamma(1+2\omega i)}{\Gamma(1+i)} \frac{[s\hat{\Xi}(s) - \Xi(0)]}{s^{2i\omega+1}} \quad (15)$$

$$\hat{L} \left[{}^{\text{GYGC}}D_x^{(\omega)}\Xi(x) \right] = \frac{1+\omega^2}{\sqrt{\pi^\omega(1-\omega)}} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega} \right)^i \frac{[s\hat{\Xi}(s) - \Xi(0)]}{\Gamma(1+i\omega) s^{2i\omega+1}} \quad (16)$$

For the details of the LT of eqs. (1), (2b), (7b), and (8b), see [2, 17, 28].

A plot of the different kernels of the FD for $\omega = 0.6$ is displayed in fig. 1.

The FD of variable order

Variable-order FD of Caputo type in term of the singular power-law kernel is given by [15, 16]:

$$D_{a^+}^{[\omega(x)]}\Xi(x) = \frac{1}{\Gamma[1-\omega(x)]} \cdot \int_a^x \frac{1}{(x-\mu)^{\omega(x)}} \left[\frac{d\Xi(\mu)}{d\mu} \right] d\mu \quad (17)$$

where $a \leq x$ and n is an integer.

Variable-order FD of Caputo-Fabrizio type in term of the exponential kernel, denoted by:

$$y = \varphi_1[x, \omega(t)] = \exp\left[-\frac{\omega(x)}{1-\omega(x)}x\right] = \sum_{i=0}^{\infty} \left[-\frac{\omega(x)}{1-\omega(x)}\right]^i \frac{x^i}{\Gamma(1+i)} \quad (18a)$$

is defined by:

$${}^{VYCF}D_x^{[\omega(x)]}\Xi(x) = \frac{1}{1-\omega(x)} \int_a^x \exp\left[-\frac{\omega(x)}{1-\omega(x)}(x-\mu)\right] \Xi^{(1)}(\mu) d\mu \quad (18b)$$

where $a \leq x$.

Variable-order FD of the stretched exponential function in the sense of the Sun-Hao-Zhang-Baleanu type, denoted by:

$$y = \varphi_2[x, \omega(t)] = \exp\left[-\frac{\omega(x)}{1-\omega(x)}x^{\omega(x)}\right] = \sum_{i=0}^{\infty} \left[-\frac{\omega(x)}{1-\omega(x)}\right]^i \frac{x^{i\omega(x)}}{\Gamma(1+i)} \quad (19a)$$

is defined by:

$${}^{VYSHZB}D_x^{(\omega)}\Xi(x) = \frac{\mathfrak{I}[\omega(x)]}{1-\omega(x)} \int_a^x \exp\left[-\frac{\omega(x)}{1-\omega(x)}(x-\mu)^{\omega(x)}\right] \Xi^{(1)}(\mu) d\mu \quad (19b)$$

where $a \leq x$.

Generalized variable-order FD of the stretched exponential function, denoted by:

$$y = \varphi_3[x, \omega(t)] = \exp[-x^{\omega(x)}] = \sum_{i=0}^{\infty} (-1)^i \frac{x^{i\omega(x)}}{\Gamma(i+1)} \quad (20a)$$

is defined:

$${}^{GVYSHZB}D_x^{(\omega)}\Xi(x) = \frac{\mathfrak{I}[\omega(x)]}{1-\omega(x)} \int_a^x \exp[-(x-\mu)^{\omega(x)}] \Xi^{(1)}(\mu) d\mu \quad (20b)$$

where $a \leq x$.

Variable-order FD involving the Mittag-Leffler function kernel, denoted by:




Figure 1. The plot of the different kernels of the FD for $\omega = 0.6$

$$y = \varphi_4[x, \omega(t)] = E_{\omega(x)} \left[-\frac{\omega(x)}{1-\omega(x)} x \right] = \sum_{i=0}^{\infty} \left[-\frac{\omega(x)}{1-\omega(x)} \right]^i \frac{x^i}{\Gamma[i\omega(x)+1]} \quad (21a)$$

is given:

$${}^{\text{VYGC}}D_x^{[\omega(x)]} \Xi(x) = \frac{1}{1-\omega(x)} \int_a^x E_{\omega(x)} \left[-\frac{\omega(x)}{1-\omega(x)} (x-\mu) \right] \Xi^{(1)}(\mu) d\mu \quad (21b)$$

Generalized variable-order FD involving the Mittag-Leffler function kernel, denoted by:

$$y = \varphi_5[x, \omega(t)] = E_{\omega(x)}(-x) = \sum_{i=0}^{\infty} (-1)^i \frac{x^i}{\Gamma[i\omega(x)+1]} \quad (22a)$$

is given:

$${}^{\text{GVYGC}}D_x^{[\omega(x)]} \Xi(x) = \frac{1}{1-\omega(x)} \int_a^x E_{\omega(x)}[-(x-\mu)] \Xi^{(1)}(\mu) d\mu \quad (22b)$$

The Atangana-Baleanu-type variable-order FD involving the stretched Mittag-Leffler function kernel, denoted by:

$$y = \varphi_6[x, \omega(t)] = E_{\omega(x)} \left[-\frac{\omega(x)}{1-\omega(x)} x^{\omega(x)} \right] = \sum_{i=0}^{\infty} \left[-\frac{\omega(x)}{1-\omega(x)} \right]^i \frac{x^{i\omega(x)}}{\Gamma[i\omega(x)+1]} \quad (23a)$$

is defined by:

$${}^{\text{VYABC}}D_x^{[\omega(x)]} \Xi(x) = \frac{1}{\Gamma[1-\omega(x)]} \int_a^x E_{\omega(x)} \left[-\frac{\omega(x)}{1-\omega(x)} (x-\mu)^{\omega(x)} \right] \Xi^{(1)}(\mu) d\mu \quad (23b)$$

where $a \leq x$.

Generalized variable-order FD involving the stretched Mittag-Leffler function kernel, denoted by:

$$y = \varphi_7[x, \omega(t)] = E_{\omega(x)}[-x^{\omega(x)}] = \sum_{i=0}^{\infty} (-1)^i \frac{x^{i\omega(x)}}{\Gamma[i\omega(x)+1]} \quad (24a)$$

is defined by:

$${}^{\text{GVYABC}}D_x^{[\omega(x)]} \Xi(x) = \frac{1}{\Gamma[1-\omega(x)]} \int_a^x E_{\omega(x)}[-(x-\mu)^{\omega(x)}] \Xi^{(1)}(\mu) d\mu \quad (24b)$$

where $a \leq x$.

The plot of the different kernels of FD for $\omega(x) = x - 1$ is displayed in fig. 2.

The LT of the function $x^{-1-\omega(x)}/\Gamma[-\omega(x)]$, given by Coimbra, takes the form [13]:

$$\hat{L} \left\{ \frac{x^{-1-\omega(x)}}{\Gamma[-\omega(x)]} \right\} = s^{\omega(x)} \quad (25)$$

The LT of the function is given by [14]:




Figure 2. The kernels of the different FD for $\omega(x) = x - 1$

$$\hat{L}\left\{E_{\omega(x)}\left[-\beta x^{\omega(x)}\right]\right\} = \frac{s^{\omega(x)-1}}{s^{\omega(x)} + \beta} \quad (26)$$

The LT of FD of variable order are:

$$\hat{L}\left\{{}^{\text{VYCF}}D_x^{[\omega(x)]}\Xi(x)\right\} = \frac{1}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{\left[-\frac{\omega(x)}{1-\omega(x)}\right]^i \Gamma(i+1) [s\hat{\Xi}(s) - \Xi(0)]}{s^{i+1}} \quad (27)$$

$$\hat{L}\left\{{}^{\text{VYSHZB}}D_x^{[\omega(x)]}\Xi(x)\right\} = \frac{\Im[\omega(x)]}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{\left[-\frac{\omega(x)}{1-\omega(x)}\right]^i \Gamma[i\omega(x)+1] [s\hat{\Xi}(s) - \Xi(0)]}{\Gamma(i+1)s^{i\omega(x)+1}} \quad (28)$$

$$\hat{L}\left\{{}^{\text{GVYSHZB}}D_x^{[\omega(x)]}\Xi(x)\right\} = \frac{\Im[\omega(x)]}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{(-1)^i \Gamma[i\omega(x)+1] [s\hat{\Xi}(s) - \Xi(0)]}{\Gamma(i+1)s^{i\omega(x)+1}} \quad (29)$$

$$\hat{L}\left\{{}^{\text{VYGC}}D_x^{[\omega(x)]}\Xi(x)\right\} = \frac{1}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{\left[-\frac{\omega(x)}{1-\omega(x)}\right]^i \Gamma(i+1) [s\hat{\Xi}(s) - \Xi(0)]}{\Gamma[i\omega(x)+1]s^{i+1}} \quad (30)$$

$$\hat{L}\left\{{}^{\text{GVYGC}}D_x^{[\omega(x)]}\Xi(x)\right\} = \frac{1}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{(-1)^i \Gamma(i+1) [s\hat{\Xi}(s) - \Xi(0)]}{\Gamma[i\omega(x)+1]s^{i+1}} \quad (31)$$

$$\hat{L}\left\{{}^{\text{VYABC}}D_x^{[\omega(x)]}\Xi(x)\right\} = \frac{1}{\Gamma[1-\omega(x)]} \frac{s^{\omega(x)-1} [s\hat{\Xi}(s) - \Xi(0)]}{s^{\omega(x)} + \frac{\omega(x)}{1-\omega(x)}} \quad (32)$$

$$\hat{L}\left\{{}^{\text{GVYABC}}D_x^{[\omega(x)]}\Xi(x)\right\} = \frac{1}{\Gamma[1-\omega(x)]} \frac{s^{\omega(x)-1} [s\hat{\Xi}(s) - \Xi(0)]}{s^{\omega(x)} + 1} \quad (33)$$

The constant- and variable-order anomalous relaxation problems arising in heat transfer

The anomalous relaxation in the sense of the constant Caputo, Caputo-Fabrizio, Sun-Hao-Zhang-Baleanu types, and variable-order FD of Caputo type were discussed in [15, 23]. To present the temperature fields in the complex media, we consider a class of anomalous relaxation equations with the aid of FD of constant and variable orders. The anomalous relaxation involving the variable-order FD are in line with the result based on the generalized versions. Therefore, we consider the LT-type solutions for the anomalous relaxation with the aid of the generalized FD of the variable order in this section.

The anomalous relaxation in the sense of FD with respect to the Mittag-Leffler function kernel reads:

$$-k {}^{YC}D_x^{(\omega)}\Xi_{YC}(x) = \Xi_{YC}(x) \quad (34a)$$

with LT of the solution:

$$\hat{\Xi}_{YC}(s) = \frac{\frac{k\mathfrak{Z}(\omega)}{1-\omega} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{\Gamma(i+1)}{\Gamma[i\omega+1]} \frac{1}{s^{i+1}}}{\frac{k\mathfrak{Z}(\omega)}{1-\omega} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{\Gamma(i+1)}{\Gamma(i\omega+1)} \frac{1}{s^i} + 1} \quad (34b)$$

where k is a constant, and $\Xi_{YC}(0) = 1$ is the initial temperature field.

The anomalous relaxation involving FD with respect to the stretched exponential function kernel in the sense of Gaussian-like type is:

$$-k {}^{GYC}D_x^{(\omega)}\Xi_{GYC}(x) = \Xi_{GYC}(x) \quad (35a)$$

with LT of the solution:

$$\hat{\Xi}_{GYC}(s) = \frac{\frac{k(1+\omega^2)}{\sqrt{\pi^\omega(1-\omega)}} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{\Gamma(1+2\omega i)}{\Gamma(1+i)} \frac{1}{s^{2i\omega+1}}}{\frac{k(1+\omega^2)}{\sqrt{\pi^\omega(1-\omega)}} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{\Gamma(1+2\omega i)}{\Gamma(1+i)} \frac{1}{s^{2i\omega}} + 1} \quad (35b)$$

where k is a constant, and $\Xi_{GYC}(0) = 1$ is the initial temperature field.

The anomalous relaxation involving FD with respect to the stretched Mittag-Leffler function kernel in the sense of Gaussian-like type can be written:

$$-k {}^{GYGC}D_x^{(\omega)}\Xi_{GYGC}(x) = \Xi_{GYGC}(x) \quad (36a)$$

with LT of the solution:

$$\hat{\Xi}_{GYGC}(s) = \frac{\frac{k(1+\omega^2)}{\sqrt{\pi^\omega(1-\omega)}} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{1}{\Gamma(1+i\omega)} \frac{1}{s^{2i\omega+1}}}{\frac{k(1+\omega^2)}{\sqrt{\pi^\omega(1-\omega)}} \sum_{i=0}^{\infty} \left(-\frac{\omega}{1-\omega}\right)^i \frac{1}{\Gamma(1+i\omega)} \frac{1}{s^{2i\omega}} + 1} \quad (36b)$$

where k is a constant, and $\Xi_{GYGC}(0) = 1$ is the initial temperature field.

The anomalous relaxation involving the variable-order FD of Caputo-Fabrizio type in term of the exponential kernel is:

$$-k^{VYCF} D_x^{[\omega(x)]} \Xi_{VYCF}(x) = \Xi_{VYCF}(x) \quad (37a)$$

with LT of the solution:

$$\hat{\Xi}_{VYCF}(s) = \frac{\frac{k}{1-\omega(x)} \sum_{i=0}^{\infty} \left[\frac{\omega(x)}{1-\omega(x)} \right]^i \frac{\Gamma(i+1)}{s^{i+1}}}{\frac{k}{1-\omega(x)} \sum_{i=0}^{\infty} \left[\frac{\omega(x)}{1-\omega(x)} \right]^i \frac{\Gamma(i+1)}{s^i} + 1} \quad (37b)$$

where k is a constant, and $\Xi_{VYCF}(0) = 1$ is the initial temperature field.

The anomalous relaxation in the sense of the variable-order FD of Caputo-Fabrizio type in term of the exponential kernel is represented:

$$-k^{GVYSHZB} D_x^{[\omega(x)]} \Xi_{GVYSHZB}(x) = \Xi_{GVYSHZB}(x) \quad (38a)$$

with LT of the solution:

$$\hat{\Xi}_{GVYSHZB}(s) = \frac{\frac{k \mathfrak{Z}[\omega(x)]}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{(-1)^i \Gamma[i\omega(x)+1]}{\Gamma(i+1) s^{i\omega(x)+1}}}{\frac{k \mathfrak{Z}[\omega(x)]}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{(-1)^i \Gamma[i\omega(x)+1]}{\Gamma(i+1) s^{i\omega(x)}} + 1} \quad (38b)$$

where k is a constant, and $\Xi_{GVYGC}(0) = 1$ is the initial temperature field.

The anomalous relaxation involving FD with respect to the stretched Mittag-Leffler function kernel in the sense of Gaussian-like type is considered:

$$-k^{GVYGC} D_x^{[\omega(x)]} \Xi_{GVYGC}(x) = \Xi_{GVYGC}(x) \quad (39a)$$

with LT of the solution:

$$\hat{\Xi}_{GVYGC}(s) = \frac{\frac{k}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{(-1)^i \Gamma(i+1)}{\Gamma[i\omega(x)+1] s^{i+1}}}{\frac{k}{1-\omega(x)} \sum_{i=0}^{\infty} \frac{(-1)^i \Gamma(i+1)}{\Gamma[i\omega(x)+1] s^i} + 1} \quad (39b)$$

where k is a constant, and $\Xi_{GVYGC}(0) = 1$ is the initial temperature field.

The anomalous relaxation in the sense of the variable-order FD of Caputo-Fabrizio type in term of the exponential kernel is represented:

$$-k^{GVYABC} D_x^{[\omega(x)]} \Xi_{GVYABC}(x) = \Xi_{GVYABC}(x) \quad (40a)$$

with LT of the solution:

$$\hat{\Xi}_{\text{GVYABC}}(s) = \frac{\frac{k}{\Gamma[1-\omega(x)]} \frac{s^{\omega(x)-1}}{s^{\omega(x)} + 1}}{\frac{k}{\Gamma[1-\omega(x)]} \frac{s^{\omega(x)}}{s^{\omega(x)} + 1} + 1} \quad (40b)$$

where k is a constant, and $\Xi_{\text{GVYABC}}(0) = 1$ is the initial temperature field.

Conclusion

In this work, we suggested a class of FD of constant and variable orders for the first time. The plots of the kernel functions of the different patterns in were discussed in detail. The LT-type solutions for the anomalous relaxations involving FD of constant and variable orders were also given. The proposed formulas are useful to open up the new prospects of describing the fractional-order heat transfer equations in the complex media.

Acknowledgement

This work is supported by the State Key Research Development Program of the People's Republic of China (Grant No.2016YFC0600705), the Natural Science Foundation of China (Grant No. 51323004), and the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD2014).

Nomenclature

k – coefficient, [m]
 x – space co-ordinate, [m]

Greek symbols

$\Xi(x)$ – temperature fields in the different FD, [K]
 ω – constant order, [–]
 $\omega(x)$ – variable order, [–]

References

- [1] West, B. J., *Fractional Calculus View of Complexity: Tomorrow's Science*, CRC Press, Boca Raton, Fla., USA, 2015
- [2] Mainardi, F., *Fractional Calculus and Waves in Linear Viscoelasticity: an Introduction to Mathematical Models*, World Scientific, Singapore, 2010
- [3] Herrmann, R., *Fractional Calculus: An Introduction for Physicists*, World Scientific, Singapore, 2014
- [4] He, J.-H., A New Fractal Derivation, *Thermal Science*, 15 (2011), Suppl. 1, pp. S145-S147
- [5] Yang, X. J., *et al.*, *Local Fractional Integral Transforms and Their Applications*, Academic Press, New York, USA, 2015
- [6] Yang, X. J., *et al.*, On Local Fractional Operators View of Computational Complexity: Diffusion and Relaxation Defined on Cantor Sets, *Thermal Science*, 20 (2016), Suppl. 3, pp. S145-S147
- [7] Yang, X. J., *et al.*, On a Fractal LC-Electric Circuit Modeled by Local Fractional Calculus, *Communications in Nonlinear Science and Numerical Simulation*, 47 (2017), June, pp. 200-206
- [8] Mainardi, F., Gorenflo, R., Time-Fractional Derivatives in Relaxation Processes: a Tutorial Survey, *Fractional Calculus and Applied Analysis*, 10 (2007), 3, pp. 269-308
- [9] Mainardi, F., Spada, G., Creep, Relaxation and Viscosity Properties for Basic Fractional Models in Rheology, *The European Physical Journal Special Topics*, 193 (2011), 1, pp. 133-160
- [10] Kilbas, A. A., *et al.*, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, The Netherlands, 2006
- [11] Samko, S. G., Ross, B., Integration and Differentiation to a Variable Fractional Order, *Integral Transforms and Special Functions*, 1 (1993), 4, pp. 277-300
- [12] Kilbas, A. A., *et al.*, *Fractional Integral and Derivatives: Theory and Applications*, Gordon and Breach, Switzerland, 1993

- [13] Lorenzo, C. F., Hartley, T. T., Variable Order and Distributed Order Fractional Operators, *Nonlinear Dynamics*, 29 (2002), 1-4, pp. 57-98
- [14] Coimbra, C. F. M., Mechanics with Variable Order Differential Operators, *Annalen der Physik*, 12 (2003), 11-12, pp. 692-703
- [15] Ramirez, L. E. S., Coimbra, C. F. M., A Variable Order Constitutive Relation for Viscoelasticity, *Annalen der Physik*, 16 (2007), 7-8, pp. 543-552
- [16] Bhrawy, A. H., Zaky, M. A., Numerical Algorithm for the Variable-Order Caputo Fractional Functional Differential Equation, *Nonlinear Dynamics*, 85 (2016), 3, pp. 1815-1823
- [17] Caputo, M., Fabrizio, M. A., New Definition of Fractional Derivative without Singular Kernel, *Progress in Fractional Differentiation and Applications*, 1 (2015), 2, pp. 73-85
- [18] Hristov, J., Transient Heat Diffusion with a Non-Singular Fading Memory: From the Cattaneo Constitutive Equation with Jeffrey's Kernel to the Caputo-Fabrizio Time-Fractional Derivative, *Thermal Science*, 20 (2016), 2, pp. 765-770
- [19] Hristov, J., Steady-State Heat Conduction in a Medium with Spatial Non-Singular Fading Memory: Derivation of Caputo-Fabrizio Space-Fractional Derivative with Jeffrey's Kernel and Analytical Solutions, *Thermal Science*, 21 (2017), 2, pp. 827-839
- [20] Atangana, A., On the New Fractional Derivative and Application to Nonlinear Fisher's Reaction - Diffusion Equation, *Applied Mathematics and Computation*, 273 (2016), Jan., pp. 948-956
- [21] Gomez-Aguilar, J. F., *et al.*, Analytical Solutions of the Electrical RLC Circuit via Liouville-Caputo Operators with Local and Non-Local Kernels, *Entropy*, 18 (2016), 8, pp. 402-413
- [22] Alkahtani, B. S. T., Atangana, A., Controlling the Wave Movement on the Surface of Shallow Water with the Caputo-Fabrizio Derivative with Fractional Order, *Chaos, Solitons & Fractals*, 89 (2016), Aug., pp. 539-546
- [23] Losada, J., Nieto, J. J., Properties of a New Fractional Derivative without Singular Kernel, *Progress in Fractional Differentiation and Applications*, 1 (2015), 2, pp. 87-92
- [24] Caputo, M., Fabrizio, M., Applications of New Time and Spatial Fractional Derivatives with Exponential Kernels, *Progress in Fractional Differentiation and Applications*, 2 (2016), 2, pp. 1-11
- [25] Yang, X.-J., *et al.*, A New Fractional Derivative without Singular Kernel: Application to the Modelling of the Steady Heat Flow, *Thermal Science*, 20 (2016), 2, pp. 753-756
- [26] Yang, A. M., *et al.*, On Steady Heat Flow Problem Involving Yang-Srivastava-Machado Fractional Derivative without Singular Kernel, *Thermal Science*, 20 (2016), Suppl. 3, pp. S717-S723
- [27] Sun, H., *et al.*, Relaxation and Diffusion Models with Non-Singular Kernels, *Physica A, Statistical Mechanics and its Applications*, 486 (2017), Feb., pp. 590-596
- [28] Atangana, A., Baleanu, D., New Fractional Derivatives with Non-Local and Non-Singular Kernel: Theory and Application to Heat Transfer Model, *Thermal Science*, 20 (2016), 2, pp. 763-769

