

THE SOLUTION OF LOCAL FRACTIONAL DIFFUSION EQUATION INVOLVING HILFER FRACTIONAL DERIVATIVE

by

*Yun QIAO, Quan-xi QIAO**

School Of Mathematics & Information Science, Henan Polytechnic University,
Jiaozuo, 454003, China

In this present work the Yang-Fourier transform method incorporating the Laplace transform method is used to solve fractional diffusion equations involving the Hilfer fractional derivative and local fractional operator. The exact solution is obtained.

Key words: *diffusion equations, Yang-Fourier transform, Laplace transform, Hilfer fractional derivative, local fractional operator*

Introduction

Nowadays, fractional calculus receives increasing attention in the scientific community. Fractional differential equation have become a useful tool in many areas of sciences and technologies. For example, fractional diffusion equation has been successfully used to describe important physical and chemical phenomena [1-6].

In recent years, the fractional diffusion equations involving the Hilfer fractional derivative and local fractional operator appear frequently in the literature[7-11]. The Hilfer fractional derivative was first proposed by Hilfer in the analysis of fractional diffusion equations, obtaining that such kind of models can be used in context of glass relaxation and aquifer problems. The theory of local fractional derivative have applied in Fokker-Planck equations, Helmholtz equations and the fractal dynamical systems etc. The main purpose of using the local fractional order derivative is to describe the anomalous diffusion phenomena in fractal media[12-14].

In this paper, we consider the following fractional order reaction-diffusion equation:

$$D_t^{\beta,\gamma} u(x,t) = \frac{\partial^{2\alpha}}{\partial x^{2\alpha}} u(x,t), \quad (1)$$

subject to the initial condition

$$J^{(1-\beta)(1-\gamma)} u(x,0) = f(x), \quad (2)$$

where $D^{\beta,\gamma}$ is a Hilfer fractional derivative of order $0 < \beta \leq 1$ and type $0 \leq \gamma \leq 1$, J^λ is

* Corresponding author ; e-mail: 13839198723@163.com

the Riemann-Liouville fractional integral of order λ , $\frac{\partial^{2\alpha}}{\partial x^{2\alpha}}$ denotes the local fractional derivative of order 2α , and $f(x)$ is a given function of x .

The rest of the paper is organized as follows. In section 2, we introduce the basic definitions and properties on the Hilfer fractional derivative and the local fractional derivative. In section 3, the Yang-Fourier transform and the Laplace transform are used to solve the problem (1)-(2) and the exact solution is obtained. Finally, the conclusions are considered in section 4.

Fractional calculus and local fractional calculus

In this section, we recall some basic definitions and properties on the Hilfer fractional derivative and the local fractional derivative which shall be used in this paper. For more detail see [8,16].

The Riemann-Liouville (R-L) fractional integral operator of order $\alpha > 0$ of a function is defined by:

$$J^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} f(s) ds, \quad (3)$$

$$J^0 f(x) = f(x). \quad (4)$$

Properties of the operator J^α can be found in [16] and we mention only the following: For $\alpha, \beta \geq 0, x > 0$ and $\lambda > -1$:

$$J^\alpha J^\beta f(x) = J^{\alpha+\beta} f(x), \quad (5)$$

$$J^\alpha J^\beta f(x) = J^\beta J^\alpha f(x), \quad (6)$$

$$J^\alpha (x^\lambda) = \frac{\Gamma(\lambda+1)}{\Gamma(1+\alpha+\lambda)} x^{\lambda+\alpha}. \quad (7)$$

The fractional derivative of $f(x)$ in Riemann-Liouville sense is defined by:

$${}^{RL}D^\mu f(y) = \frac{d^n}{dy^n} J^{n-\mu} f(y), \quad n = [\text{Re}(\mu)] + 1. \quad (8)$$

The Caputo fractional derivative is defined by:

$${}^C D^\mu f(y) = J^{n-\mu} f^{(n)}(y). \quad (9)$$

We recall the following basic properties:

$${}^C D^\alpha J^\alpha f(x) = f(x), \quad (10)$$

$$J^{\alpha c} D^{\alpha} f(x) = f(x) - \sum_{k=0}^{m-1} f^{(k)}(0^+) \frac{x^k}{k!}, \quad x > 0 \quad (11)$$

Hilfer generalized the fractional derivative (8) by the following fractional derivative of order $0 < \beta \leq 1$ and type $0 \leq \gamma \leq 1$:

$$D^{\beta, \gamma} f(y) = J^{\gamma(1-\beta)} \frac{d}{dy} J^{(1-\gamma)(1-\beta)} f(y), \quad (12)$$

which is called Hilfer fractional derivative. Note that when $0 < \beta \leq 1, \gamma = 0$, the Hilfer fractional derivative would be the R-L fractional derivative:

$${}^{RL}D^{\beta} f(y) = \frac{d}{dy} J^{1-\beta} f(y), \quad (13)$$

when $0 < \beta \leq 1, \gamma = 1$, it corresponds to the Caputo fractional derivative:

$${}^cD^{\beta} f(y) = J^{1-\beta} f'(y), \quad (14)$$

On the Laplace transform of the Hilfer fractional derivative (12), in Ref.[8], it is found for $0 < \beta < 1$ that

$$L[D^{\beta, \gamma} f(y)] = s^{\beta} L[f(y)] - s^{\gamma(\beta-1)} J^{(1-\beta)(1-\gamma)} f(0),$$

where the initial value term

$$J^{(1-\beta)(1-\gamma)} f(0), \quad (15)$$

is evaluated in the limit $y \rightarrow 0^+$, in the space of the Lebesgue integrable functions

$$L[0, \infty] = \left\{ f : \|f\|_1 = \int_0^{\infty} |f(y)| dy < \infty \right\}.$$

Next, we recall the notations and some properties of the local fractional operators[16-17].

The function $f(x)$ is called the local fractional continuous (LFC) at $x = x_0$, if it is valid for

$$|f(x) - f(x_0)| < \varepsilon^{\alpha}, \quad 0 < \alpha \leq 1 \quad (16)$$

with $|x - x_0| < \delta$, for $\varepsilon > 0$. If $f(x)$ is the LFC for any $x \in (a, b)$, then it is called LFC on the interval (a, b) , denoted by $f(x) \in C_{\alpha}(a, b)$.

Setting $f(x) \in C_{\alpha}(a, b)$, local fractional derivative (LFD) of $f(x)$ at $x = x_0$ is defined as

$$\frac{d^\alpha}{dx^\alpha} f(x)|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Delta^\alpha (f(x) - f(x_0))}{(x - x_0)^\alpha}, \quad (17)$$

where $\Delta^\alpha (f(x) - f(x_0)) \cong \Gamma(1 + \alpha)\Delta(f(x) - f(x_0))$.

The local fractional integral (LFI) of $f(x)$ in the interval $[a, b]$ is defined

$$\int_a^b f(t)(dt)^\alpha = \lim_{\Delta t \rightarrow 0} \sum_{j=0}^{N-1} f(t_j)(\Delta t_j)^\alpha, \quad (18)$$

where $\{t_j | j = 0, 1, \dots, N\}$ is a partition of the interval $[a, b]$, $\Delta t_j = t_j - t_{j-1}$, $\Delta t = \{\Delta t_j\}$, $t_0 = a$, and $t_N = b$.

If $f(x) \in C_\alpha(a, b)$, we define its Yang-Fourier transform:

$$F_\alpha[f(x); \omega] = f_\omega^{F, \alpha}(\omega) = \frac{1}{\Gamma(1 + \alpha)} \int_{-\infty}^{+\infty} E_\alpha(-i^\alpha \omega^\alpha x^\alpha) f(x)(dx)^\alpha \quad (19)$$

and its inverse Yang-Fourier transform

$$f(x) = F_\alpha^{-1}[f_\alpha^{F, \alpha}(\omega)] = \frac{1}{(2\pi)^\alpha} \int_{-\infty}^{+\infty} E_\alpha(i^\alpha \omega^\alpha x^\alpha) f_\alpha^{F, \alpha}(\omega)(d\omega)^\alpha. \quad (20)$$

Here, $E_\alpha(z)$ is the Mittag-Leffer function:

$$E_\alpha(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + 1)}. \quad (21)$$

We also use the following two-parameter Mittag-Leffer function in the next section:

$$E_{\alpha, \beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}. \quad (22)$$

The following formulas are valid:

$$F_\alpha[af(x) + bg(x); \omega] = aF_\alpha[f(x); \omega] + bF_\alpha[g(x); \omega], \quad (23)$$

$$F_\alpha\left[\frac{d^\alpha}{dx^\alpha} f(x); \omega\right] = i^\alpha \omega^\alpha F[f(x); \omega]; \quad (24)$$

$$L[t^{(\mu-1)(1-\nu)} E_{\mu, 1-(1-\mu)(1-\nu)}(-\lambda t^\mu)] = \frac{s^{-\nu(1-\mu)}}{s^\mu + \lambda}. \quad (25)$$

The solution of the problem (1)-(2)

In this section, we investigate the problem (1)-(2). Our main result is as follows :

The solution of the problem (1)-(2) is given by

$$u(x,t) = \int_{-\infty}^{+\infty} f(\xi)v(x-\xi,t)(d\xi)^\alpha, \quad (24)$$

where

$$v(x,t) = F_\alpha^{-1}[g(\omega,t);x] \quad (25)$$

and

$$g(\omega,t) = t^{(\beta-1)(1-\gamma)} E_{\beta,\beta-\gamma(\beta-1)}(-t^\beta i^\alpha \omega^\alpha). \quad (26)$$

Let $u^*(\omega,t)$ denotes the Yang-Fourier transform of the function $u(x,t)$ with respect to x . Taking the Yang-Fourier transform of both sides of equation (1) yields:

$$D_t^{\beta,\gamma} u^*(\omega,t) = i^\alpha \omega^\alpha u^*(\omega,t). \quad (27)$$

Applying the Laplace transform to (27), we obtain:

$$s^\beta L[u^*(\omega,t);s] - s^{\gamma(\beta-1)} F_\alpha[f(x);\omega] = i^\alpha \omega^\alpha L[u^*(\omega,t);s].$$

So

$$L[u^*(\omega,t);s] = \frac{s^{\gamma(\beta-1)} f^*(\omega)}{s^\beta - i^\alpha \omega^\alpha}, \quad (28)$$

where $f^*(\omega) = F_\alpha[f(x);\omega]$.

From the inverse Laplace transform of the relation (28), and the (25), we get:

$$u^*(\omega,t) = f^*(\omega) t^{(\beta-1)(1-\gamma)} E_{\beta,\beta-\gamma(\beta-1)}(t^\beta i^\alpha \omega^\alpha). \quad (29)$$

Let

$$t^{(\beta-1)(1-\gamma)} E_{\beta,\beta-\gamma(\beta-1)}(t^\beta i^\alpha \omega^\alpha) = g(\omega,t),$$

and

$$F_\alpha^{-1}[g(\omega,t);x] = v(x,t).$$

Then, by using the convolution theorem of the Yang-Fourier transform, we get

$$u(x,t) = \int_{-\infty}^{+\infty} f(\xi)v(x-\xi,t)(d\xi)^\alpha. \quad (30)$$

This completes the proof.

For $0 < \beta \leq 1, \gamma = 1$, the problem (1)-(2) becomes

$$\begin{cases} {}^c D_t^\beta u(x,t) = \frac{\partial^{2\alpha}}{\partial x^{2\alpha}} u(x,t), \\ u(x,0) = f(x). \end{cases}$$

Then the solution is

$$u(x,t) = \int_{-\infty}^{+\infty} f(\xi)v(x-\xi,t)(d\xi)^\alpha,$$

where

$$v(x,t) = F_\alpha^{-1}[E_\beta(t^\beta i^\alpha \omega^\alpha); x].$$

For $0 < \beta \leq 1, \gamma = 0$, the problem (1)-(2) becomes

$$\begin{cases} {}^{RL}D_t^\beta u(x,t) = \frac{\partial^{2\alpha}}{\partial x^{2\alpha}} u(x,t), \\ J^{1-\beta} u(x,0) = f(x). \end{cases}$$

Then the solution is

$$u(x,t) = \int_{-\infty}^{+\infty} f(\xi)v(x-\xi,t)(d\xi)^\alpha,$$

where

$$v(x,t) = F_\alpha^{-1}[t^{\beta-1} E_{\beta,\beta}(t^\beta i^\alpha \omega^\alpha); x].$$

Conclusion

In this paper, the Yang-Fourier transform and the Laplace transform have been used to solve a new fractional diffusion equations involving the Hilfer fractional derivative and local fractional operator and the exact solution is obtained.

Acknowledgements

This paper was partially supported by Henan Natural Science Foundation in China under Grant No. 182300410105, and by the Doctoral Foundation at Henan Polytechnic University in China under Grant No. B2015-52.

References

- [1] Mainardi, Francesco, Y. Luchko, and G. Pagnini. The fundamental solution of the space-time fractional diffusion equation, *Fractional Calculus and Applied Analysis*, 2(2007), 4, pp.153--192
- [2] Langlands, T. A. M., and B. I. Henry, The accuracy and stability of an implicit solution method for the fractional diffusion equation, *Journal of Computational Physics*, 2(2005), 205, pp. 719-736
- [3] Tadjerana, Charles, A second-order accurate numerical approximation for the fractional diffusion equation, *Journal of Computational Physics*, 1(2006), 213, pp. 205-213
- [4] X.-J. Yang, F. Gao, Y. Ju., *General Fractional Derivatives with Applications in*

Viscoelasticity, New York: Academic Press, 2019

- [5] Yang, X.-J. *General Fractional Derivatives: Theory, Methods and Applications*, New York: CRC Press, 2019
- [6] Saadatmandi, Abbas, and M. Dehghan, A tau approach for solution of the space fractional diffusion equation, *Computers and Mathematics with Applications* , 3(2011), 62, pp. 1135-1142
- [7] Tomovski, et al. Fractional and operational calculus with generalized fractional derivative operators and Mittag–Leffler type functions, *Integral Transforms & Special Functions*, 11(2010), 21, pp. 797-814
- [8] Katugampola, Udit N, A New Approach to Generalized Fractional Derivatives, *Mathematics* , 4(2011), 6, pp. 1--15
- [9] Sandev, Trifce, R. Metzler, and Živorad Tomovski, Fractional diffusion equation with a generalized Riemann–Liouville time fractional derivative, *Journal of Physics A Mathematical & Theoretical* , 25(2011), 44, pp. 255203
- [10] Zivorad Tomovski, Generalized Cauchy type problems for nonlinear fractional differential equations with composite fractional derivative operator, *Nonlinear Analysis Theory Methods and Applications*, 7(2012), 75, pp.3364-3384
- [11] Qassim, M. D, K. M. Furati, and N. E. Tatar, On a Differential Equation Involving Hilfer-Hadamard Fractional Derivative, *Abstract and Applied Analysis* , 1085-3375(2014) , pp. 1-9
- [12] Yang, X. J., et al., *Local Fractional Integral Transforms and Their Applications*. New York: Academic Press, 2015
- [13] Saxena, Ram K., Z. Tomovski, and T. Sandev, Fractional Helmholtz and fractional wave equations with Riesz-Feller and generalized Riemann-Liouville fractional derivatives, *Eprint Arxiv* , 3(2014), 7, pp.24-32
- [14] Yang, Xiao Jun, D. Baleanu, and H. M. Srivastava, Local fractional similarity solution for the diffusion equation defined on Cantor sets, *Applied Mathematics Letters*, 47(2015), 1, pp. 54-60
- [15] Yang, X. J. *Advanced Local Fractional Calculus and Its Applications*. New York: World Science Publisher, 2012
- [16] Oldham, Keith B, and J. Spanier, The fractional calculus, *Mathematical Gazette* , 247(1974), 56, pp.396-400
- [17] Yang, X. J. *Local Fractional Functional Analysis & Its Applications*. Hong Kong: Asian Academic Publisher, 2011

Paper submitted: April 21, 2018

Paper revised: July 19, 2018

Paper accepted: October 11, 2018