

INVESTIGATING THE IMPACTS OF AIR QUALITY AND WEATHER INDICATORS ON THE SPREAD OF SARS-CoV-2 IN ISTANBUL, TURKEY

by

Caner TANIS^a and Kadir KARAKAYA^{b*}

^a Faculty of Science, Cankiri Karatekin University, Cankiri, Turkey

^b Faculty of Science, Selcuk University, Konya, Turkey

Original scientific paper

<https://doi.org/10.2298/TSCI201126071T>

Air pollution is having a positive impact on the spread of the SARS-CoV-2 virus. The effects of meteorological parameters on the spread of SARS-CoV-2 are a matter of curiosity. The main purpose of this paper is to determine the association between air quality indexes ($PM_{2.5}$, PM_{10} , NO_2 , SO_2 , CO , and O_3) and weather parameters (temperature, humidity, pressure, dew, and wind speed) with the number of SARS-CoV-2 cases, hospitalizations, hospital discharges. In this paper, we also focused on determining the impact of air pollution and weather parameters on the number of daily hospitalizations and daily discharges. It is gleaned daily cases, hospitalizations, hospital discharges, meteorological, and air quality data in Istanbul from Turkey between July 15, 2020, and September 30, 2020. We performed the Pearson correlation analysis to evaluate the effects of meteorological parameters and air quality indexes on the variables related to SARS-CoV-2. It is determined a statistically significant positive relationship between air quality indexes such as CO , SO_2 , $PM_{2.5}$, PM_{10} , NO_2 , and the number of daily confirmed SARS-CoV-2 cases. We also observed a negative association between weather parameters such as temperature and pressure and the number of daily confirmed SARS-CoV-2 cases. Our study proposes that high air quality could reduce the number of SARS-CoV-2 cases. The empirical findings of this paper might provide key input to prevent the spread of SARS-CoV-2 across Turkey.

Key words: *air quality index, SARS-CoV-2, weather parameters*

Introduction

The new Coronavirus (SARS-CoV-2) firstly appearing in Wuhan, China, in December 2019, and rapidly has crossed borders, infecting people throughout the whole world. As of October 6, 2020, it is globally reported that there have been 35 347 404 confirmed cases of SARS-CoV-2 with 1 039 406 deaths by the World Health Organization [1]. The SARS-CoV-2 is officially announced as a global pandemic on March 11, 2020 [2]. The SARS-CoV-2 is essentially contaminated human-to-human *via* some factors such as transmitted surfaces, close contact, and respiratory droplets. The contamination of SARS-CoV-2 is closely similar to other respiratory viruses such as influenza [2-7]. The contagion of viruses may be affected by environmental and meteorological factors such as temperature, humidity, air quality, population density, and use of masks [8].

* Corresponding author, e-mail: istatistikadir@gmail.com

Recently, many studies in the literature illustrated that air quality, temperature and population density are correlated with the number of SARS-CoV-2 cases. Coskun *et al.* [9] mentioned that SARS-CoV-2 virus in the air is a threatening factor for humans in Turkey. Sahin [10] investigated the number of SARS-CoV-2 case relationships with the parameters of weather such as humidity, temperature, and wind speed for eleven cities in Turkey. They reported that negative impacts of the average temperature, humidity, and dew point on the approved SARS-CoV-2 cases number. Iddrisu *et al.* [11] emphasized that the number of daily confirmed SARS-CoV-2 cases is negatively related to maximum temperature and humidity while it is positively correlated with rainfall and maximum wind. Moreover, it is determined maximum temperature, rainfall, and humidity were significant predictors of daily confirmed cases of SARS-CoV-2 in Ghana [11]. It is investigated the relationship between air pollution levels and the risk of SARS-CoV-2 in Italy by [12]. It is analyzed the effects of weather indicators on the spread of SARS-CoV-2 in Spain by [13]. Investigation of the relationship between temperature and SARS-CoV-2 in China is examined by [14]. Malki *et al.* [15] examined the relationship between the parameters of weather and the rate of mortality due to SARS-CoV-2. They demonstrated that the weather parameters are more relevant in predicting the mortality rate of SARS-CoV-2 virus when compared to the other demographic variables such as population, age, and urbanization. It was also emphasized that there is a negative relationship between the number of SARS-CoV-2 cases in some countries, including the USA, Italy [15]. The main climate factor that suppresses the spread of SARS-CoV-2 is high solar radiation. High temperatures and wind speed also are potential factors to prevent the spread of SARS-CoV-2 [16]. In [17] evaluated the relationship between environmental parameters and the growth rate of SARS-CoV-2. It is expressed that the spread rate of the global pandemic is related to the changes in temperature and found to be the most effective environmental factor that could control the spread by new cases/day with a 1 °C rise in it. The most effective temperature range for the spread of SARS-CoV-2 is -6.28 °C and 14.51 °C [17].

Stieb *et al.* [18] found that there is a positive relationship between SARS-CoV-2 incidence and long-term PM_{2.5} exposure. Chauhan and Singh [19] reported a decline in PM_{2.5} concentration of major cities (New York, Los Angeles, Rome, Zaragoza, Dubai, Mumbai, Shanghai, and Beijing) due to less mobility of the population and keep social distancing in order to prevent the spread of SARS-CoV-2. The low PM_{2.5} concentrations reflect high air quality and the efforts made in the major cities to control the spread of infection of SARS-CoV-2 [19]. Badr *et al.* [20] constructed a mathematical model between mobility patterns and transmission of SARS-CoV-2 in the USA. Bontempi [21] concluded that it is not conspicuous direct associations between the existence of a high level of PM₁₀ and the dispersal of the SARS-CoV-2 for Lombardy (Italy).

Yar *et al.* [22] investigated the spatial correlation between PM₁₀, PM_{2.5}, and SARS-CoV-2 deaths in China and concluded that a higher concentration of PM₁₀ and PM_{2.5} positively correlated with deaths due to SARS-CoV-2 virus. On the other hand, the growth of SARS-CoV-2 pandemic was not correlated with rainfall, humidity, wind speed, NO, NO₂, O₃, and PM_{2.5}, while it was significantly associated with daily temperature in Japan [3]. Sarkodie and Owusu [23] reported that high temperature and high RH reduce the activity, continuity, and contamination of SARS-CoV-2 for 20 countries including, Turkey, Italy, Russia, Netherlands, US, Spain, Portugal, France, Germany, United Kingdom, Belgium, India, Iran, China, Brazil, Ecuador, Canada, Switzerland, Peru, and Saudi Arabia. Currently, Turkey is one of the highly affected countries with approximately 329 000 total confirmed cases and more than 8000 deaths [24]. Coskun *et al.* [9] emphasized that population density positively affects the spread

of SARS-CoV-2 in Turkey. Baser [25] observed that population density is a significant predictor of infection rates of SARS-CoV-2 virus. Istanbul is the most crowded city in Turkey, with approximately 15.5 million citizens, one of Europe's most important metropolises. Therefore, we preferred to study on Istanbul.

The previous studies provided that meteorological variables and air quality are correlated with the spread of SARS-CoV-2 pandemic. Coskun *et al.* [9] emphasized the availability of the positive relationship between population density and SARS-CoV-2 cases. Therefore, we prefer to analyze Istanbul, which is the most populous city in Turkey. The motivate of this research is the lack of a special study on the association between weather parameters and the spread of the SARS-CoV-2 virus and its relation to air pollution for Istanbul in the literature. In this regard, this study focuses on to determine the association between the confirmed SARS-CoV-2 cases, hospitalizations, hospital discharges, and weather parameters such as temperature, humidity, pressure, dew, wind speed, and also examine its correlation between air quality indexes such as PM_{2.5}, PM₁₀, O₃, NO₂, SO₂, CO in Istanbul, Turkey. This study aims to assess the impacts of weather conditions and air pollution on the spread of SARS-CoV-2 in Istanbul, Turkey.

Materials and methods

Study design and areas

This research was designed to evaluate the relationship between the growth of SARS-CoV-2 pandemic and the environmental parameters during the period of the pandemic. The increasing number of confirmed SARS-CoV-2 cases led to various measures to be taken to control the spread of the SARS-CoV-2 pandemic by the Turkish government. Some of these precautions in Turkey can be listed: It was temporarily restricted to entry or exit abroad. It was declared curfews in metropolitan provinces on the weekends. Education was temporarily suspended at all levels. The number of passengers in public transportation was limited. Some places, such as malls, restaurants, cafes, were temporarily closed. The flexible period of working began in public institutions. Other countries around the world also applied similar restrictions as Turkey. It is discussed some restrictions worldwide and international laws in detail by [26].

On the other hand, social activities have also been affected by SARS-CoV-2 pandemic. Some organizations such as conferences, theater, cinema, concert, and wedding were banned until a second announcement. National football leagues were postponed. One of the most crowded metropolises in Europe is Istanbul, which is essentially affected by SARS-CoV-2 pandemic. Istanbul is Turkey's most populous city, mostly topped the list of the number of cases SARS-CoV-2 in Turkey. We motivated for this study due to a lack of research on the relation between air pollution and some variables related to SARS-CoV-2, such as the number of cases, the number of hospitalizations, and discharges due to SARS-CoV-2 for Istanbul province. Thus, Istanbul is chosen as the area of study due to it is the most affected province by SARS-CoV-2, and the most crowded city in Turkey.

In this regard, it is considered the analysis period from July 15 to September 30, 2020, to evaluate the correlation of the pandemic growth with the environmental and meteorological factors.

The main aim of this study is to determine the association between weather parameters and air quality and the spread of SARS-CoV-2 pandemic. Unlike other investigations, this paper provides the relationship of air quality and weather parameters with some parameters

regarding SARS-CoV-2, such as the number of cases, hospitalizations, and discharges from the hospital rather than an only association between the number of SARS-CoV-2 cases and air pollution or weather parameters. The temperature [°C], humidity [%], air pressure, dew point [°C], and wind speed [kmh] are chosen as meteorological indicators. Further, PM_{2.5}, PM₁₀, O₃, NO₂, SO₂, and CO are determined as air quality indexes to assess air pollution of the province.

Data collection

We consider analyzing variables regarding weather and air quality into four periods. The data of these variables are collected based on four time-frames, namely on the day of the case, within 3, 7, and 14 days of the case since the incubation period of SARS-CoV-2 virus varies from 1 day to 14 days. The data related to SARS-CoV-2 cases, hospitalizations, and discharges from the hospital were collected from <https://covid19.saglik.gov.tr>. Daily air quality and meteorological measurement data were obtained from [27]. In this research, it is used the maximum available data of all variables from 15th July to 30th September 2020.

Statistical analysis

Pearson correlation coefficient is a statistic that measures linear correlation between two variables. In this study, Pearson correlation is used to determine the correlation between air quality indexes or meteorological indicators and variables regarding SARS-CoV-2 (cases, hospitalizations, and discharges from the hospitals). Pearson coefficient correlation is estimated using the following formula:

$$r_{\text{pearson}} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}}$$

where n is sample of size. The statistical tests are applied using R software, and the significance level of 0.05 was considered.

Results

The descriptive results are given in tab. 1. From tab. 1, it is observed that the daily number of cases varied between 99 and 268 from July, 15 to September, 30 in Istanbul.

Table 2 provides the results of pearson correlation tests conducted to determine the relationship between air quality index and the spread of SARS-CoV-2. Firstly, the air index PM₁₀ is the most correlated with the number of daily cases. A statistically significant positive relationship was observed between the number of cases on that day and the PM₁₀ observed 7 days ago ($r = 0.532$). The air index having the second-highest correlation with the daily number of cases is PM_{2.5}. A positive correlation was determined between the PM_{2.5} value observed 14 days ago and the number of daily cases ($r = 0.517$). It can be concluded that the high degree of PM₁₀ and PM_{2.5} allows the spread of SARS-CoV-2. The height of PM₁₀ and PM_{2.5}, two of the most important indicators of air pollution, was associated with the increasing number of SARS-CoV-2 cases. Moreover, O₃ is the air quality index that has the highest negative relationship with the daily number of cases. It is determined the negative correlation between the daily number of cases and O₃ observed on the day ($r = -0.469$). It is observed that the air quality indexes observed 7 days ago generally are more related to the number of

Table 1. Descriptive statistics

| Variables | <i>n</i> | Mean | Std. dev. | Min | Max | <i>p</i> ₅ | <i>p</i> ₉₅ | Skewness | Kurtosis |
|-----------------------------------|----------|-----------|-----------|-------|-------|-----------------------|------------------------|----------|----------|
| SARS-CoV-2 variables | | | | | | | | | |
| Daily case | 78 | 183.9487 | 4.0311 | 99 | 268 | 117.7 | 249.1 | -0.069 | -0.790 |
| Number of new hospitalizations | 78 | 57.8462 | 15.7067 | 27 | 123 | 31.95 | 82.10 | 0.787 | 2.770 |
| Number of new hospital discharges | 78 | 62.1026 | 24.5703 | 25 | 114 | 30.9 | 110.0 | 0.541 | -0.872 |
| Air quality index | | | | | | | | | |
| CO | 78 | 126.4679 | 139.9968 | 58.60 | 500.0 | 60.2650 | 500.0 | 2.184 | 3.181 |
| SO ₂ | 78 | 95.4141 | 130.2914 | 39.1 | 500 | 40.395 | 500.0 | 2.751 | 6.039 |
| PM ₁₀ | 78 | 188.4872 | 301.7965 | 44 | 999 | 47.95 | 999 | 2.301 | 3.525 |
| PM _{2.5} | 77 | 60.9740 | 18.8092 | 30 | 116 | 36.6 | 17 | 1.164 | 1.368 |
| O ₃ | 78 | 87.6705 | 77.8564 | 27.30 | 476.7 | 42.9 | 287.12 | 3.258 | 11.487 |
| NO ₂ | 78 | 98.6705 | 128.9549 | 24.5 | 500 | 31.26 | 500 | 2.759 | 6.125 |
| Weather parameters | | | | | | | | | |
| Temperature | 78 | 29.4064 | 2.4457 | 20 | 34.5 | 24.975 | 33 | -1.057 | 2.638 |
| Dew | 78 | 20.0128 | 1.5373 | 16 | 23 | 17 | 22 | -0.644 | -0.121 |
| Humidity | 78 | 91.3679 | 4.6570 | 79.5 | 100 | 82 | 100 | -0.388 | -0.254 |
| Pressure | 78 | 1014.8128 | 3.2266 | 1008 | 1022 | 1009.9 | 1021 | 0.144 | -0.317 |
| Wind speed | 78 | 8.3667 | 1.8438 | 3.30 | 14.10 | 3.79 | 11.35 | -0.259 | 1.888 |

daily cases compared to the others. This situation may have occurred due to the incubation period of the SARS-CoV-2 virus. Secondly, O₃ has the highest correlation with the number of hospitalized patients. The O₃ value observed 3 days ago having a negatively effect on the daily number of hospitalized patients ($r = -0.283$). The CO is one of the two air quality indices associated with the daily number of hospitalized patients. There is a negative correlation between the daily number of hospitalized patients and the CO value observed 7 days ago ($r = -0.246$), tab. 2. Thirdly, O₃ observed 3 days ago highest positively correlated with the daily number of discharges from the hospital among air quality indexes ($r = 0.429$). The PM_{2.5} has the highest negative relationship with the daily number of people discharged from the hospital. The PM_{2.5} observed on that day has a statistically significant association with the daily number of discharges ($r = -0.385$). Thus, high O₃ has a positive influence on the number of daily discharges, while high PM_{2.5} has a negative effect.

When the relationship between weather indicators and the spread of SARS-CoV-2 is evaluated according to tab. 3. First of all, the pressure is the most associated with the number of daily cases. The pressure observed 7 days ago statistically significant positive relationship with the number of cases ($r = 0.592$). Maximum temperature observed 7 days ago highest negatively correlated the number of cases on the day ($r = -0.552$). Furthermore, it is determined a statistically significant relationship between maximum humidity and the number of daily cases, tab. 3. The maximum humidity observed 3 days ago has the highest positive cor-

relation with the number of daily hospitalizations ($r = 0.317$). Secondly, it can be concluded that there is no significant association between the maximum temperature and the number of hospitalizations. Lastly, it is clearly seen that maximum pressure negatively affected the number of discharges from the hospital. Thus, the weather parameters such as temperature, dew, and humidity have a negative effect on the number of the daily case while the pressure is positively affected. On the other hand, there is no relationship between wind speed and the spread of SARS-CoV-2.

Table 2. Correlation analysis between air quality index and SARS-CoV-2

| | Air quality index | Daily case | Number of new hospitalizations | Number of new hospital discharges |
|-------------|-------------------|------------|--------------------------------|-----------------------------------|
| On the day | CO | 0.302** | 0.140 | -0.285* |
| | SO ₂ | 0.235* | 0.009 | -0.254* |
| | PM ₁₀ | 0.344** | -0.002 | -0.263* |
| | PM _{2.5} | 0.439** | -0.093 | -0.385** |
| | O ₃ | -0.469** | -0.268* | 0.303** |
| | NO ₂ | 0.278* | -0.027 | -0.250* |
| 3 days ago | CO | 0.338** | -0.183 | -0.181 |
| | SO ₂ | 0.275* | -0.132 | -0.208 |
| | PM ₁₀ | 0.325** | -0.154 | -0.229* |
| | PM _{2.5} | 0.365** | -0.135 | -0.384** |
| | O ₃ | -0.405** | -0.283* | 0.429** |
| | NO ₂ | 0.312** | -0.182 | -0.255* |
| 7 days ago | CO | 0.394** | -0.246* | -0.231* |
| | SO ₂ | 0.421** | -0.158 | -0.183 |
| | PM ₁₀ | 0.532** | -0.107 | -0.099 |
| | PM _{2.5} | 0.330** | -0.055 | -0.325** |
| | O ₃ | -0.288* | -0.031 | 0.167 |
| | NO ₂ | 0.417** | -0.170 | -0.201 |
| 14 days ago | CO | 0.194 | -0.009 | -0.148 |
| | SO ₂ | 0.331** | 0.014 | -0.107 |
| | PM ₁₀ | 0.276* | 0.175 | -0.007 |
| | PM _{2.5} | 0.517* | -0.028 | -0.132 |
| | O ₃ | -0.135 | -0.054 | -0.003 |
| | NO ₂ | -0.285* | -0.039 | -0.145 |

* Significance at the 5% level, ** significance at the 1% level.

When tab. 2 is investigated, one of the first striking features is that the highest relationships generally appeared before 7 days. Also, when tabs. 2 and 3 are examined, it is clearly seen that the relationship between the number of cases and observed variables 14 days ago

Table 3. Correlation analysis between meteorological parameters and SARS-CoV-2

| | Weather indicators | Daily case | Number of new hospitalizations | Number of new hospital discharges |
|-------------|--------------------|------------|--------------------------------|-----------------------------------|
| On the day | Temperature | -0.402** | 0.058 | 0.190 |
| | Dew | -0.195 | -0.228* | -0.018 |
| | Humidity | -0.038 | 0.309** | 0.193 |
| | Pressure | 0.243* | -0.055 | -0.290* |
| | Wind Speed | -0.180 | 0.103 | 0.057 |
| 3 days ago | Temperature | -0.518** | -0.076 | -0.031 |
| | Dew | -0.377** | -0.026 | -0.007 |
| | Humidity | -0.166 | 0.317** | 0.088 |
| | Pressure | 0.309** | 0.011 | -0.386** |
| | Wind Speed | -0.080 | 0.053 | 0.218 |
| 7 days ago | Temperature | -0.552** | -0.112 | 0.130 |
| | Dew | -0.344** | -0.190 | -0.044 |
| | Humidity | -0.256* | 0.135 | 0.089 |
| | Pressure | 0.592** | 0.233* | -0.133 |
| | Wind Speed | 0.107 | 0.054 | 0.074 |
| 14 days ago | Temperature | -0.139 | -0.103 | 0.067 |
| | Dew | 0.060 | 0.063 | 0.123 |
| | Humidity | -0.106 | 0.111 | 0.204 |
| | Pressure | 0.257* | -0.098 | 0.097 |
| | Wind Speed | 0.205 | 0.185 | 0.214 |

* Significance at the 5% level, ** significance at the 1% level.

is less than variables measured at other time intervals. Although CO, SO₂, PM₁₀, PM_{2.5}, and NO₂ generally have a positive relationship with the daily case. There is a negative relationship with the number of new hospital discharges. In addition, no statistically significant relationship was found between many variables and the number of new hospitalizations. The O₃ is known as a natural disinfectant that does not leave any residue. In this study, it is also determined that O₃ has a negative relationship with the daily case. However, it has a positive relationship with the number of new hospital discharges. This feature of O₃ can be a good research subject for future studies. From tab. 2, it is also seen that the highest relationship is determined between PM₁₀ and daily case for before 7 days.

From tab. 3, it can be seen that the temperature is a negative relationship with the daily case. The highest relationship between daily case and temperature is determined 7 days ago. There is no relationship between temperature and number of new hospitalizations, and number of new hospital discharges. Observed maximum dew 3, 7 days ago is negatively affected the daily cases. Humidity has a positive relationship with the number of new hospitali-

zations for the same day and before 3 days. From tab. 3, highest positive relationship is found between daily case and pressure observed 7 days ago. Also, the pressure measured on the day and 3 days ago has a negative effect on the number of new hospital discharges. In tab. 3, it is seen that there is no relationship between wind speed and all weather parameters for all situations.

Discussion

The aim of this study is to determine the effects of air pollution and weather parameters on the spread of SARS-CoV-2. Unlike previous studies, which aimed to only assess the correlation between air pollution or meteorological indicators and SARS-CoV-2 cases, this paper provides the effects of weather parameters and air quality indexes on some parameters related to SARS-CoV-2, such as confirmed cases, hospitalizations, discharges from the hospital due to SARS-CoV-2.

Scientists emphasized that moisture is an important factor in preventing the spread of SARS-CoV-2 [28]. In the literature, most investigations (33 out of 61) suggest a negative relationship between SARS-CoV-2 and temperature in various countries such as the US, Japan, Ghana, Spain, Italy, China. Similarly, many papers (13 out of 27) present a negative impact of humidity worldwide. Another study shows that other weather parameters such as wind speed, rainfall, and radiation clearly did not relate to SARS-CoV-2 [29]. Rosario *et al.* [16] discussed the association between weather parameters (temperature, humidity, solar radiation, wind speed, and precipitation) and SARS-CoV-2 infection in the State of Rio de Janeiro, Brazil. They concluded that solar radiation strongly negatively correlated with the incidence of SARS-CoV-2. Prata *et al.* [30] indicated that tropical temperatures had a negative linear correlation with the number of confirmed SARS-CoV-2 cases in Brazil. Dogan *et al.* [8] investigated the impacts of meteorological parameters on SARS-CoV-2 in New Jersey, USA. They found maximum temperature negatively correlated, while humidity and air quality, $PM_{2.5}$, positively affect the spread of SARS-CoV-2 in New Jersey. It was observed the higher is the number of SARS-CoV-2 cases, the higher average wind speed in 14 days, on the other hand, it was concluded that the lower the temperature on a day, the higher is the number of SARS-CoV-2 cases on that day. Moreover, the higher number of SARS-CoV-2 cases is closely related to the crowd in a city of Turkey [10]. Coskun *et al.* [9] reported that population, population density and wind speed are positive correlated with the number of SARS-CoV-2 cases for 81 of Turkey cities.

On the other hand, previous studies have determined that respiratory infectious diseases are correlated with environmental air pollutants [31-35]. Bashir *et al.* [36] found that there is a statistically significant association between environmental factors (PM_{10} , $PM_{2.5}$, SO_2 , CO, and NO_2) and SARS-CoV-2 cases and deaths in California. Environmental pollutants are significantly associated with the number of SARS-CoV-2 cases and deaths in California [36]. Cao *et al.* [31] and Yongjian *et al.* [35] reported positive correlations of $PM_{2.5}$, PM_{10} , NO_2 , and O_3 with the number of SARS-CoV-2 cases, however, a negative relationship was determined for SO_2 for China. The results of many studies [8, 18, 19, 22] indicate $PM_{2.5}$ and PM_{10} positively affect the spread of SARS-CoV-2. These results are providing evidence that air quality and weather parameters such as temperature and humidity are actually important to control the spread of SARS-CoV-2 pandemic.

Our findings supported previous studies. First of all, it is observed the highest positive correlation between before 7 days maximum PM_{10} and daily SARS-CoV-2 cases among air quality indexes ($r = 0.532$).

It is well-known that O₃ has a very high oxidation power. It is observed before 3 days maximum O₃ has the most negative effect on the number of daily new hospitalizations ($r = -0.283$). Further, it is clearly seen that examined all air quality indexes are closely related to the number of new discharges from the hospital due to SARS-CoV-2. Secondly, it is observed many statistically significant correlation between meteorological indicators and SARS-CoV-2 parameters such as cases, hospitalizations, discharges from the hospital. Before 7 days, maximum temperature has one of the most negative impact on the number of confirmed SARS-CoV-2 cases in meteorological indicators ($r = -0.552$). Before 3 days, maximum pressure has the highest level of negative effect on the number of discharges from the hospital due to SARS-CoV-2 ($r = -0.386$). Before 7 days, maximum humidity negatively affects the number of SARS-CoV-2 cases ($r = -0.256$), and wind speed clearly does not affect the number of SARS-CoV-2 cases as the results of previous studies, tab. 3. As it can be clearly seen from tab. 1, this may be due to the low wind speed (approximately 12.87 kmh on average). Our study provides new evidence that air pollution and weather parameters affects the spread of SARS-CoV-2 pandemic. Unlike the previous study, one of the important advantages of this study is that the association between air pollutants or weather parameters and SARS-CoV-2 pandemic is evaluated in many ways.

Conclusion

In conclusion, we are motivated to study the impacts of weather parameters and air quality on the number of SARS-CoV-2 cases, hospitalizations, and discharges from the hospital. We present new evidence that air quality indexes such as maximum CO, SO₂, PM_{2.5}, PM₁₀, and NO₂ have a positive effect on the number the SARS-CoV-2 cases, whereas O₃ negatively affects it. In weather parameters, maximum temperature, dew point, and humidity negatively correlated the number of SARS-CoV-2 cases while pressure positively related to it. On the other hand, it is not determined a statistically significant correlation between wind speed and SARS-CoV-2 cases. It is concluded that examined weather parameters have an impact on the spread of SARS-CoV-2 within 7 days. It is generally observed that weather parameters before 14 days have not an impact on SARS-CoV-2 parameters. We recommend the authors, who will study the relationship between the spread of SARS-CoV-2 and weather parameters, to analyze the meteorological indicators 7 days ago and the SARS-CoV-2 data of that day.

References

- [1] ***, WHO, 2020a, <https://covid19.who.int/>, Access date: October 7, 2020
- [2] ***, WHO, 2020b, WHO Director-General's Opening Remarks at the Media Briefing on SARS-CoV-2 - 11 March 2020, World Health Organization, Geneva
- [3] Azuma, K., *et al.*, Impact of Climate and Ambient Air Pollution on the Epidemic Growth During SARS-CoV-2 Outbreak in Japan, *Environmental Research*, 2020 (2020), Nov., ID 110042
- [4] Cheng, V. C. C., *et al.*, Escalating Infection Control Response to the Rapidly Evolving Epidemiology of the Coronavirus Disease 2019 (SARS-CoV-2) Due to SARS-CoV-2 in Hong Kong, *Infect. Control Hosp. Epidemiol.*, 41 (2020), 5, pp. 493-498
- [5] Lai, C. C., *et al.*, Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) and Coronavirus Disease-2019 (SARS-CoV-2): the Epidemic and the Challenges, *Int. J. Antimicrob.*, 55 (2020), 3, ID 105924
- [6] Sungnak, W., *et al.*, SARS-CoV-2 Entry Factors are Highly Expressed in Nasal Epithelial Cells Together with Innate Immune Genes, *Nat. Med.*, 26 (2020), 5, pp. 681-687
- [7] ***, WHO, 2020c Transmission of SARS-CoV-2: Implications for Infection Prevention Precautions, Scientific Brief on 9 July 2020, World health Organization, Geneva

- [8] Dogan, B., et al., Investigating the Effects of Meteorological Parameters on SARS-CoV-2: Case Study of New Jersey, United States, *Environmental Research*, 191 (2020), Dec., 110148
- [9] Coskun, H., et al., The Spread of SARS-CoV-2 Virus Through Population Density and Wind in Turkey Cities, *Science of The Total Environment*, 751 (2021), Jan., 141663
- [10] Sahin, M., Impact of weather on SARS-CoV-2 pandemic in Turkey, *Science of The Total Environment*, 728 (2020), Aug., ID 138810
- [11] Iddrisu, W. A., et al., Effects of Weather and Policy Intervention on SARS-CoV-2 Infection in Ghana 2020, On-line first, arXiv Preprint arXiv:2005.00106, 2020
- [12] Fattorini, D., Regoli, F., Role of the Chronic Air Pollution Levels in the Covid-19 Outbreak Risk in Italy, *Environmental Pollution*, 264 (2020), Sept., ID 114732
- [13] Shahzad, K., et al., Effects of Climatological Parameters on the Outbreak Spread of COVID-19 in Highly Affected Regions of Spain, *Environ. Sci. Pollut. Res. Int.*, 27 (2020), 31, pp. 39657-39666
- [14] Shahzad, F., et al., Asymmetric Nexus Between Temperature and COVID-19 in the Top Ten Affected Provinces of China: A current Application of Quantile-on-Quantile Approach, *Science of The Total Environment*, 736 (2020), Sept., ID 139115
- [15] Malki, Z., et al., Association Between Weather Data and SARS-CoV-2 Pandemic Predicting Mortality Rate: Machine Learning Approaches, *Chaos, Solitons & Fractals*, 138 (2020), Sept., ID 110137
- [16] Rosario, D. K. A., et al., Relationship Between SARS-CoV-2 and Weather: Case Study in a Tropical Country, *International Journal of Hygiene and Environmental Health*, 229 (2020), Aug., ID 113587
- [17] Sil, A., Kumar, V. N., Does Weather Affect the Growth Rate of SARS-CoV-2, A Study to Comprehend Transmission Dynamics on Human Health, *Journal of Safety Science and Resilience*, 1 (2020), 1, pp. 3-11
- [18] Stieb, D. M., et al., An Ecological Analysis of Long-Term Exposure to PM_{2.5} and Incidence of SARS-CoV-2 in Canadian Health Regions, *Environmental research*, 191 (2020), Dec., ID 110052
- [19] Chauhan, A., Singh, R. P., Decline in PM_{2.5} Concentrations Over Major Cities Around the World Associated with SARS-CoV-2, *Environmental Research*, 187 (2020), Aug., ID 109634
- [20] Badr, H. S., et al., Association Between Mobility Patterns and COVID-19 Transmission in the USA: A Mathematical Modelling Study, *The Lancet Infectious Diseases*, 20 (2020), 11, pp. 1247-1254
- [21] Bontempi, E., First Data Analysis About Possible SARS-CoV-2 Virus Airborne Diffusion Due to Air Particulate Matter (PM): The Case of Lombardy (Italy), *Environmental Research*, 186 (2020), July, ID 109639
- [22] Yao, Y., et al., Spatial Correlation of Particulate Matter Pollution and Death Rate of SARS-CoV-2, *medRxiv*, (2020)
- [23] Sarkodie, S. A., Owusu, P. A., Impact of Meteorological Factors on SARS-CoV-2 Pandemic: Evidence from Top 20 Countries with Confirmed Cases, *Environmental Research*, 191 (2020), Dec., ID 110101
- [24] ***, Republic of Turkey Ministry of Health., 2020. <https://covid19.saglik.gov.tr/>, Access date: October 7, 2020
- [25] Baser, O., Population Density Index and Its Use for Distribution of Covid-19: A Case Study Using Turkish Data, *Health Policy*, 125 (2020), 2, pp. 148-154
- [26] Ní Ghráinne, B., Covid-19, Border Closures, and International Law, (2020)
- [27] ***, <https://aqicn.org/data-platform/covid19/>, Access date: October 7, 2020
- [28] ***, <https://www.cnnturk.com/turkiye/bilimsel-arastirma-covid-19un-yayilimini-azaltmak-icin-nem-onemli>, Access date: December 27, 2020
- [29] Briz-Redón, Á., Serrano-Aroca, Á., The Effect of Climate on the Spread of the SARS-CoV-2 Pandemic: A Review of Findings, and Statistical and Modelling Techniques, *Progress in Physical Geography: Earth and Environment*, 44 (2020), 5, pp. 591-604
- [30] Prata, D. N., et al., Temperature Significantly Changes SARS-CoV-2 Transmission in (Sub) Tropical Cities of Brazil, *Science of the Total Environment*, 729 (2020), Aug., ID 138862
- [31] Cao, H., et al., Associations of Ambient Air Pollutants and Meteorological Factors with SARS-CoV-2 Transmission in 31 Chinese Provinces: A Time-Series Study, *medRxiv*, On-line first, <https://doi.org/10.1101/2020.06.24.20138867>, 2020
- [32] Gehring, U., et al., Air Pollution and the Development of Asthma from Birth Until Young Adulthood, *The European respiratory journal*, 56 (2020), 1, ID 2000147
- [33] Kan, H. D., et al., Relationship Between Ambient Air Pollution and Daily Mortality of Sars in Beijing, *Biomedical and Environmental Sciences*, 18 (2005), 1, pp. 1-4

- [34] Liu, X. X., *et al.*, Effects of Air Pollutants on Occurrences of Influenza-Like Illness and Laboratory-Confirmed Influenza in Hefei, China, *International journal of biometeorology*, 63 (2019), 1, pp. 51-60
- [35] Yongjian, Z., *et al.*, Association Between Short-Term Exposure to Air Pollution and SARS-CoV-2 Infection: Evidence from China, *Science of the total environment*, 727 (2020), July, 138704
- [36] Bashir, M. F., *et al.*, Correlation Between Environmental Pollution Indicators and SARS-CoV-2 Pandemic: A Brief Study in Californian context, *Environmental Research*, 187 (2020), Aug., 109652