

**CHEMICAL AND THERMODYNAMIC PROPERTIES
OF *BOMBYX MORI* – DOMESTIC SILK MOTH
Empirical Formula, Driving Force, and Biosynthesis,
Catabolism and Metabolism Reactions**

by

**Marko E. POPOVIĆ^{a,*}, Marijana R. PANTOVIĆ PAVLOVIĆ^{a,b},
and Gavriilo M. ŠEKULARAC^a**

^a Institute of Chemistry, Technology and Metallurgy, University of Belgrade,
Belgrade, Serbia

^b Centre of Excellence in Chemistry and Environmental Engineering – ICTM,
University of Belgrade, Belgrade, Serbia

Original scientific paper
<https://doi.org/10.2298/TSCI230901242P>

*Biothermodynamics is a discipline which has developed intensely during the last 50 years. Thermodynamic properties have been reported for humans, animals, plants, and microorganisms. However, this paper reports for the first time the empirical formula and thermodynamic properties for insects. Thermodynamic properties can be applied in research on thermodynamic interactions between organisms and their environment, as well as between organisms themselves. This paper reports for the first time the empirical formula and reactions of catabolism, biosynthesis and entire metabolism for *Bombyx mori* (domestic silk moth), as well as the thermodynamic properties of *Bombyx mori*. It is shown that growth of *Bombyx mori* is tightly related to catabolism of carbohydrates and lipids, which represents the driving force for the entire metabolism.*

Key words: insect, Gibbs energy, growth, bioenergetics, enthalpy, entropy

Introduction

The root of biothermodynamics goes back to the late 18th century in the works of the father and son Carnot [1-3], as well as Lavoisier and Laplace [4, 5]. Since then, thermodynamics has crystalized as one of the best based scientific disciplines. Some of the founders of thermodynamics are also the founders of the sub-discipline of biothermodynamics. Lavoisier and Laplace developed the first calorimeter and one of their first measurements was the metabolic heat of a Guinea pig [4, 5]. The mathematical and philosophical basis of thermodynamics was laid in the mid-19th century by Clausius [6-8]. The work of Clausius represents the basis of engineering thermodynamics [9, 10], which has developed very rapidly in the late 19th and the first half of 20th century. The development of biothermodynamics has been much slower, until the second half of the 20th century. The interest in biothermodynamics was raised by Schrodinger in his famous lecture series and book entitled *What is Life?* [11]. Prigogine has gone a step further with the development of non-equilibrium thermodynamics, as an excellent tool for analysis of processes performed by living organisms [10, 12-19]. An important step for development of

* Corresponding author, e-mail: marko.popovic@ihtm.bg.ac.rs

biothermodynamics was given by Morowitz [20-26]. During the late 20th century, Hansen has analyzed the change in entropy during life processes, as well as applications of calorimetry and thermodynamics in analysis of metabolic processes [27-38]. Von Stockar has named Gibbs energy of growth as the driving force for growth of microorganisms, and made great contributions to applications of thermodynamics and calorimetry in analysis of metabolism and in bioengineering [39-48]. Battley has developed tools for thermodynamic characterization of microorganisms [49-59]. An important research on thermodynamic characterization of bacteriophages and other microorganisms was made by Maskow [60-70]. Thermodynamic background of microorganism evolution was reported by Hansen [27-29] and Skene [71]. The application of thermodynamics in soil science was intensely developed by Barros [72-76]. Lucia has applied thermodynamics to analyze topics in life sciences, which include virus-host interactions and virus epidemiology [77-84]. The development of biothermodynamics of organisms at different hierarchical levels of organization was made by Ozilgen [85-91]. Thermodynamic characterization of organisms was reported by Popovic [92-101].

In the literature, chemical and thermodynamic properties were reported for multicellular organisms, human tissues [101], animals, plants [101], bacteria [95, 99] and viruses [77-79, 84, 91-94, 96-98, 102-107]. However, until now insects have not yet been a subject of research of biothermodynamics. Thus, chemical and thermodynamic characterization of insects has not yet been performed.

The molecular composition of *Bombyx mori* larva was reported by Morowitz [25]. *Bombyx mori*, the domestic silk moth, belongs to the moth family (*Bombycidae*). Its closest relative is the wild silk moth (*Bombyx mandarina*). The larva of *Bombyx mori* is the silkworm. It has been very important for millennia, since it is used to produce silk. *Bombyx mori* larva feed on the leaves of mulberry (*Morus alba*), which are rich in protein [108].

Bombyx mori represents a biothermodynamic system, which contains a highly organized amount of substance, clearly separated from its environment. It exchanges matter and energy with its environment and exhibits growth, which appears as a consequence of organization and accumulation of matter. Thus, *Bombyx mori* performs catabolic and anabolic processes.

The goal of this paper is to perform chemical and thermodynamic characterization of *Bombyx mori*. Moreover, the goal is to determine thermodynamic properties of catabolism, anabolism and entire metabolism, as well as to formulate catabolism, biosynthesis and total metabolism reactions.

Methods

Data sources

Molecular composition of *Bombyx mori* larvae was taken from [25]. The molecular composition of *Bombyx mori* larvae is 55.5% protein, 13.3% lipid and 1.8% carbohydrate (in mass fractions) [25].

The empirical formulas of protein and lipids was taken from [109]. The empirical formula of protein is $\text{CH}_{1.59}\text{O}_{0.32}\text{N}_{0.26}\text{S}_{0.007}$ (or $\text{C}_{100}\text{H}_{159}\text{N}_{26}\text{O}_{32}\text{S}_{0.7}$) [109]. The empirical formula of lipids is $\text{CH}_{1.92}\text{O}_{0.12}$ (or $\text{C}_{51}\text{H}_{98}\text{O}_6$) [109]. The empirical formula of carbohydrates is CH_2O [39, 40]. Standard specific enthalpies of combustion of proteins, lipids and carbohydrates were taken from [110]. Standard specific enthalpy of combustion of protein is 22.2 MJ/kg [110]. Standard specific enthalpy of combustion of lipids is 39.8 MJ/kg [110]. Standard specific enthalpy of combustion of lipids is 18.0 MJ/kg [110].

Biosynthesis rate and metabolic heat rate of *Bombyx mori* were taken from [111, 112], respectively. The biosynthesis rate of *Bombyx mori* larvae is $r_{bs} = 2.53 \cdot 10^{-6}$ C-mol/C-mol s

(153%wt./week) [111]. The metabolic heat rate of *Bombyx mori* is $\dot{q}_{\text{met}} = -1.49 \cdot 10^{-4}$ kW/s [112].

Empirical formulas

Elemental composition of live matter was calculated using the molecular composition method [113]. The molecular composition method uses macromolecular composition of live matter as input [113]. The macromolecular composition comes in form of mass fractions of the macromolecular components of live matter [113]. These are first converted into mole fractions of macromolecular components:

$$x_X = \frac{\frac{w_X}{M_r(X)}}{\sum_Y \frac{w_Y}{M_r(Y)}} \quad (1)$$

where x_X is the mole fraction of macromolecular component X , w_X and w_Y are the mass fractions of macromolecular components X and Y , respectively, while $M_r(X)$ and $M_r(Y)$ are the molar masses of empirical formulas of molecular components X and Y , respectively [113]. The summation is over all Y macromolecular components that constitute live matter [113]. The mole fractions of macromolecular components are used to find the empirical formula of live matter, using:

$$n_J = \sum_X x_X n_J(X) \quad (2)$$

where n_J is the number of atoms of element J in live matter and $n_J(X)$ – the number of atoms of element J in the empirical formula of macromolecular component X [113]. The summation is over all X macromolecular components that constitute the live matter [113].

Thermodynamic properties of live matter

Empirical formulas were used to predict standard thermodynamic properties of live matter, through the Patel-Erickson model [52, 114], Battley model [50, 59] and Hurst-Harrison model [115, 116]. The Patel-Erickson model was used to find enthalpy, while the Battley model was used to find entropy. Enthalpy and entropy were combined to find Gibbs energy. Heat capacity was calculated through the Hurst-Harrison model.

Enthalpy of live matter

The Patel-Erickson model gives standard enthalpy of combustion of live matter, based on its empirical formula [52, 114]:


$$\Delta_C H_m^0 = -111.14 \frac{\text{kJ}}{\text{C-mol}} E \quad (3)$$

where $\Delta_C H_m^0$ is standard enthalpy of combustion of live matter and E is the number of electrons transferred to oxygen during oxidation [52, 114]. The E can be found from the equation:

$$E = 4n_C + n_H - 2n_O - 0n_N + 5n_P + 6n_S \quad (4)$$

where n_J is the number of atoms of element J in the empirical formula of live matter [52, 114].

The $\Delta_C H_m^0$ is then converted into standard enthalpy of formation, $\Delta_f H_m^0$, of live matter, through the Hess's law [52, 95]. The $\Delta_C H_m^0$ is defined as the enthalpy change for the reaction of complete oxidation of organic matter by oxygen:



Standard enthalpies of formation of all the inorganic compounds in this reaction are known [117, 118]. Thus, the Hess's law can be applied to calculate $\Delta_f H_m^0$, of live matter, through [95, 117, 118]:

$$\Delta_f H_m^0(\text{bio}) = n_C \Delta_f H^0(\text{CO}_2) + \frac{n_H}{2} \Delta_f H^0(\text{H}_2\text{O}) +$$

$$+ \frac{n_P}{4} \Delta_f H^0(\text{P}_4\text{O}_{10}) + n_S \Delta_f H^0(\text{SO}_3) - \Delta_C H_m^0 \quad (6)$$

Entropy of live matter

The Battley model is a predictive biothermodynamic model that can be used to find standard molar entropy of live matter, S_m^0 [50, 59]:

$$S_m^0(\text{bio}) = 0.187 \sum_J \frac{S_m^0(J)}{a_J} n_J \quad (7)$$

where $S_m^0(\text{bio})$ is standard molar entropy of live matter, $S_m^0(J)$ – the standard molar entropy of element J in its standard state, a_J – the number of atoms of element J in its standard state, and n_J – the number of atoms of element J in the empirical formula of live matter [50, 59]. The summation is over all J elements constituting the live matter [50, 59]. For example, the standard state of hydrogen is H_2 , with an entropy of 130.684 J/mol K and $a_H = 2$ [117].

The Battley model can also give standard entropy of formation, $\Delta_f S_m^0$, of live matter [50, 95]. In that case it takes the form:

$$\Delta_f S_m^0(\text{bio}) = -0.813 \sum_J \frac{S_m^0(J)}{a_J} n_J \quad (8)$$

where $\Delta_f S_m^0$ is then combined with $\Delta_f H_m^0$ to find standard Gibbs energy of formation, $\Delta_f G_m^0$, of live matter, using:

$$\Delta_f G_m^0 = \Delta_f H_m^0 - T \Delta_f S_m^0 \quad (9)$$

where T is temperature.

Heat capacity of live matter

Heat capacity of live matter can be found from its empirical formula through the Hurst-Harrison model [115, 116]. The Hurst-Harrison model gives standard molar heat capacity at constant pressure, $C_{p,m}^0$ for live matter:

$$C_{p,m}^0(\text{bio}) = \sum_J c_J n_J \quad (10)$$

where c_J is the contribution of element J to the heat capacity, while n_J – the number of atoms of element J in the empirical formula of live matter [115, 116]. The values of c_J for different elements can be found in [115].

Specific thermodynamic properties of live matter

Thermodynamic properties of live matter discussed previously are all in molar form (per C-mole of live matter). Sometimes, it is more convenient to have them expressed in specific form (per gram of live matter). Thus, standard thermodynamic properties of live matter were also reported in specific form. Specific thermodynamic properties can be calculated from molar properties:

$$X_g = \frac{X_m}{M_r} \quad (11)$$

where X_g is the property in specific form, X_m – the corresponding property in molar form, and M_r – the molar mass of live matter. This equation was applied to $\Delta_f H^0$, S^0 , $\Delta_f G^0$, C_p^0 , $\Delta_c H^0$, and $\Delta_f S^0$.

Biosynthesis, catabolism and metabolism reactions

Based on empirical formulas, macrochemical equations were constructed that describe biosynthesis, catabolism and metabolism. These reactions, which describe the overall stoichiometry of major metabolic processes, have been applied with great success in bioengineering [39, 40, 43, 45], microbiology [49, 51, 52, 57, 58, 92, 93, 102, 103], plant science [100] and soil research [74].

Biosynthesis reactions

Biosynthesis reactions are macromolecular equations that describe conversion of nutrients by organisms into new live matter [39, 40, 49, 119]. They are sometimes called anabolic reactions [39, 40, 119]. Each biosynthesis reaction consists of nutrients as reactants, live matter as the main product of biosynthesis and additional biosynthesis products. Every nutrient taken by the organism from the environment contributes one or more elements to the formation of new live matter [120]. The biosynthesis reaction for *Bombyx mori* has the general form:




where ν_{pr} , $\nu_{\text{H}_2\text{O}}$, ν_{CO_2} , ν_{bio} , ν_{NH_3} , and ν_{O_2} represent the stoichiometric coefficients of proteins, water, CO_2 , live matter, ammonia, and oxygen, respectively. In this work the convention with positive stoichiometric coefficients for products and negative stoichiometric coefficients for reactants was used [117, 118]. The $\text{CH}_{1.59}\text{O}_{0.32}\text{N}_{0.26}$ is the empirical formula of proteins [109] and $\text{C}_{nc}\text{H}_{nh}\text{O}_{no}\text{N}_{nn}$ is the empirical formula of live matter.

Proteins are the source of carbon and nitrogen. Water is a source of hydrogen atoms needed for production of live matter. The main product is new live matter with an empirical formula $\text{C}_{nc}\text{H}_{nh}\text{O}_{no}\text{N}_{nn}$. Excess nitrogen is released as urea ($\text{CH}_4\text{N}_2\text{O}$), which is the end product of nitrogen metabolism. The O_2 takes excess electrons produced by oxidation of nutrients. It will disappear from the product side, when the catabolic reaction is added.

Catabolism reactions

Catabolism reactions describe degradation of nutrients by organisms into simple catabolic products, to provide energy that drives the metabolism [39, 40, 49, 119]. A catabolism reaction contains nutrients and the electron acceptor as reactants, and on the products side catabolic waste products [39, 40, 49]. The general catabolism reaction for *Bombyx mori* has the form:




where v_{lp} , v_{ch} , v_{O_2} , v_{H_2O} and v_{CO_2} , are the stoichiometric coefficients of lipids, carbohydrates, oxygen, water and carbon dioxide, respectively. No new live matter is produced in the catabolism reaction [39, 40, 49]. The $CH_{1.92}O_{0.12}$ is the empirical formula of lipids [109] and CH_2O is empirical formula of carbohydrates. Lipids and carbohydrates represent the main sources of energy. The ratio of lipids and carbohydrates in *Morus alba* (mulberry) leaves, which represent the main food for *Bombyx mori* larvae [111], was taken from [108]. Oxygen is the electron acceptor, which takes electrons from the nutrients to release energy [40]. The catabolic waste products are CO_2 and H_2O . The CO_2 takes the oxidized carbon atoms. The H_2O takes excess hydrogen.

Metabolism reactions

Biosynthesis and catabolism are the main components of metabolism [40, 119]. Thus, the biosynthesis and catabolism reactions are added to obtain the metabolism reaction. The catabolism reaction is multiplied with the reciprocal of the biomass yield, Y (the method for finding Y will be considered below) and then added to the biosynthesis reaction [40, 49]. Thus, the stoichiometric coefficient of molecule X in the metabolism reaction, $v_{X,met}$, can be found:

$$v_{X,met} = \frac{1}{Y} v_{X,cat} + v_{X,bs} \quad (14)$$

where Y is the biomass yield, while $v_{X,cat}$ and $v_{X,bs}$ are the stoichiometric coefficients of molecule X in the catabolism and biosynthesis reactions, respectively [40]. Thus, the metabolism reaction has the general form:



Thermodynamic properties of biosynthesis, catabolism and metabolism

Once the biosynthesis, catabolism and metabolism reactions and stoichiometric coefficients are known, their thermodynamic properties can be determined. This is done through the Hess's law [117, 118]. Standard thermodynamic properties of reaction r are given by:

$$\Delta_r H^0 = \sum_X v_X \Delta_f H^0(X) \quad (16)$$

$$\Delta_r S^0 = \sum_X v_X S_m^0(X) \quad (17)$$

$$\Delta_r G^0 = \sum_X v_X \Delta_f G^0(X) \quad (18)$$

$$\Delta_r C_{p,m}^0 = \sum_X v_X C_{p,m}^0(X) \quad (19)$$

where $\Delta_r H^0$, $\Delta_r S^0$, $\Delta_r G^0$, and $\Delta_r C_{p,m}^0$ are standard enthalpy, entropy, Gibbs energy and heat capacity of reaction r , respectively, v_X is the stoichiometric coefficient of substance X in reaction r , $\Delta_f H^0(X)$ standard enthalpy of formation of substance X , $S_m^0(X)$ standard molar entropy of substance X , $\Delta_f G^0(X)$ standard Gibbs energy of substance X , and $C_{p,m}^0(X)$ standard molar heat capacity at constant pressure of substance X [117, 118]. The summation is over all X substances that react in reaction r [117, 118]. The convention with positive stoichiometric coefficients for products and negative stoichiometric coefficients for reactants was used [117, 118]. Except

for thermodynamic parameters, this work also analyzes the kinetic parameters of metabolism, which are described in the next subsection.

Kinetic properties of metabolism

Kinetic properties of metabolism determined in this research include biosynthesis heat rate, \dot{q}_{bs} , catabolism heat rate, \dot{q}_{cat} , catabolism rate, r_{cat} , and biomass yield, Y . The input are literature values of total metabolic heat rate, \dot{q}_{met} , and biosynthesis rate, r_{bs} . In the first step, the standard enthalpy of biosynthesis, $\Delta_{bs}H^0$, was multiplied with the biosynthesis rate, r_{bs} , to find the biosynthesis heat rate, \dot{q}_{bs} [110, 121, 122]:

$$\dot{q}_{bs} = r_{bs} \cdot \Delta_{bs}H^0 \quad (20)$$

Biosynthesis and catabolism are the two main components of metabolism [39, 40, 119]. Thus, according to the Hess's law [117, 118], the metabolic heat rate, \dot{q}_{met} , is the sum of the biosynthesis heat rate and catabolism heat rate, \dot{q}_{cat} :

$$\dot{q}_{met} = \dot{q}_{bs} + \dot{q}_{cat} \quad (21)$$

This means that the catabolic heat rate is:

$$\dot{q}_{cat} = \dot{q}_{met} - \dot{q}_{bs} \quad (22)$$

This equation was used to find \dot{q}_{cat} .

The catabolism heat rate was then used to find the catabolism reaction rate. Catabolism reaction rate can be found by dividing the catabolism heat rate, \dot{q}_{cat} , with standard enthalpy of catabolism, $\Delta_{cat}H^0$ [110, 121, 122]:

$$r_{cat} = \frac{\dot{q}_{cat}}{\Delta_{cat}H^0} \quad (23)$$

Finally, rates of biosynthesis and catabolism were used to find the biomass yield. The ratio of r_{bs} and r_{cat} is equal to the biomass yield, Y [40]:

$$Y = \frac{r_{bs}}{r_{cat}} \quad (24)$$

The biomass yield Y is then used to combine the biosynthesis and catabolism reactions into the metabolism reaction with eq. (14), as described previously.

Phenomenological equations and coefficients

The kinetic and thermodynamic parameters are united through the framework of non-equilibrium thermodynamics. This is done by using phenomenological equations. Phenomenological equations give rates of processes based on their physical driving forces [39, 110, 123]. The physical driving force of chemical reactions is Gibbs energy [39, 110, 123]. Gibbs energy of a chemical reaction can be used to find its affinity [123]. This is done through:

$$A_i = -\frac{\Delta_i G}{T} \quad (25)$$

where T is temperature, while A_i and $\Delta_i G$ are affinity and Gibbs energy of reaction i , respectively [123]. This equation was used to find biosynthesis affinity, A_{bs} , and catabolism affinity, A_{cat} , based on their Gibbs energies.

If a system performs multiple chemical reactions simultaneously, the rate of every reaction can be found through the phenomenological equation:

$$r_i = \sum_j L_{ij} A_j \quad (26)$$

where r_i is the rate of reaction i , L_{ij} the phenomenological coefficient between reactions i and j , while A_j is affinity of reaction j [123]. The summation is over all j chemical reactions that the system performs [123].

In our case, we have a system with two reactions – biosynthesis and catabolism. Thus, the phenomenological equations take the form:

$$r_{\text{bs}} = L_{\text{BB}}A_{\text{bs}} + L_{\text{BC}}A_{\text{cat}} \quad (27)$$

$$r_{\text{cat}} = L_{\text{BC}}A_{\text{bs}} + L_{\text{CC}}A_{\text{cat}} \quad (28)$$

where r_{bs} and r_{cat} are the rates of the biosynthesis and catabolism reactions, respectively, while A_{bs} and A_{cat} the affinities (driving forces) of biosynthesis and catabolism reactions, respectively [39, 123]. The L_{BB} , L_{CC} , and L_{BC} are the phenomenological coefficients [39, 123]. The L_{BB} describes the influence of the biosynthesis driving force on biosynthesis rate [39, 123]. The L_{CC} describes the influence of catabolism driving force on catabolism rate [39, 123]. The L_{BC} is called the coupling coefficient, since it describes the influence of catabolism on biosynthesis and *vs.* [39, 123].

Under complete coupling between biosynthesis and catabolism, the coupling coefficient can be found [39]:

$$L_{\text{BC}} = \sqrt{L_{\text{BB}}L_{\text{CC}}} \quad (29)$$

Finally, when eqs. (27)-(29) can be combined and solved for L_{BB} and L_{CC} . This gives:

$$L_{\text{BB}} = \frac{r_{\text{bs}}^2}{r_{\text{cat}}A_{\text{cat}} + r_{\text{bs}}A_{\text{bs}}} \quad (30)$$

$$L_{\text{CC}} = \frac{r_{\text{cat}}^2}{r_{\text{cat}}A_{\text{cat}} + r_{\text{bs}}A_{\text{bs}}} \quad (31)$$

Equations (29)-(31) were used to find the L_{BB} , L_{CC} , and L_{BC} .

Results

This paper reports for the first time the empirical formula of an insect – *Bombyx mori* (silkworm). It was determined through the molecular composition method [113]. It is given in tab. 1. The empirical formula of the *Bombyx mori* larva is $\text{CH}_{1.6803}\text{O}_{0.2815}\text{N}_{0.1903}\text{S}_{0.005122}$, which has a molar mass of 21.04 g/C-mol.

Table 1. Empirical formula and molar mass of live matter of *Bombyx mori* (silkworm) larva; the empirical formula has the general form $\text{C}_{n_{\text{C}}}\text{H}_{n_{\text{H}}}\text{O}_{n_{\text{O}}}\text{N}_{n_{\text{N}}}\text{S}_{n_{\text{S}}}$

n_{C}	n_{H}	n_{O}	n_{N}	n_{S}	Mr [gC-mol ⁻¹]
1	1.6803	0.2815	0.1903	0.005122	21.04

Thermodynamic properties of live matter are reported for the first time in this paper for an insect. They are given in tab. 2. They were determined based on the empirical formula, through the Patel-Erickson [52, 114] and Battley models [50, 59]. These include standard enthalpy of formation, $\Delta_f H^\circ$, standard molar entropy, S_m° , standard Gibbs energy of formation, $\Delta_f G^\circ$, standard molar heat capacity at constant pressure, $C_{p,m}^\circ$, standard enthalpy of combustion, $\Delta_c H^\circ$, and standard entropy of formation, $\Delta_f S^\circ$. All the properties were reported per C-mole, X_m , and per gram, X_g . For the live matter of *Bombyx mori* larva, standard enthalpy of formation is -63.52 kJ/C-mol (-3.020 kJ/g), standard molar entropy is 30.40 J/C-molK (1.445 J/gK),


standard Gibbs energy of formation is -24.12 kJ/C-mol (-1.146 kJ/g), standard molar heat capacity at constant pressure is 31.00 J/C-mol K (1.474 J/gK), standard enthalpy of combustion is -572.16 kJ/C-mol (-27.197 kJ/g) and standard entropy of formation is -132.17 J/C-molK (-6.283 J/gK).

Table 2. Thermodynamic properties of live matter of *Bombyx mori* (silkworm) larva: standard enthalpy of formation, $\Delta_f H^0$, standard molar entropy, S_m^0 , standard Gibbs energy of formation, $\Delta_f G^0$, standard molar heat capacity at constant pressure, $C_{p,m}^0$, standard enthalpy of combustion, $\Delta_c H^0$, standard entropy of formation, $\Delta_f S^0$; all the properties are given per C-mole, X_m , and per gram, X_g


Per C-mole	$\Delta_f H_m^0$ [kJC-mol ⁻¹]	S_m^0 [JC-mol ⁻¹ K ⁻¹]	$\Delta_f G_m^0$ [kJC-mol ⁻¹]	$C_{p,m}^0$ [JC-mol ⁻¹ K ⁻¹]	$\Delta_c H_m^0$ [kJC-mol ⁻¹]	$\Delta_f S_m^0$ [JC-mol ⁻¹ K ⁻¹]
	-63.52	30.40	-24.12	31.00	-572.16	-132.17

Per gram	$\Delta_f H_g^0$ [kJg ⁻¹]	S_g^0 [Jg ⁻¹ K ⁻¹]	$\Delta_f G_g^0$ [kJg ⁻¹]	$C_{p,g}^0$ [Jg ⁻¹ K ⁻¹]	$\Delta_c H_g^0$ [kJg ⁻¹]	$\Delta_f S_g^0$ [Jg ⁻¹ K ⁻¹]
	-3.020	1.445	-1.146	1.474	-27.197	-6.283

Biosynthesis, catabolism and entire metabolism reactions were formulated for the first time for an insect, based on the empirical formula. They are given in tab. 3. The biosynthesis reaction for the *Bombyx mori* larva:



where $\text{CH}_{1.59}\text{O}_{0.32}\text{N}_{0.26}$ is the empirical formula of protein, $\text{CH}_{1.6803}\text{O}_{0.2815}\text{N}_{0.1903}$ is the empirical formula of *Bombyx mori* live matter, tab. 1, and $\text{CH}_4\text{N}_2\text{O}$ is the formula of urea. The catabolic reaction for *Bombyx mori* larva:



where $\text{CH}_{1.92}\text{O}_{0.12}$ is the empirical formula of lipids and CH_2O the empirical formula of carbohydrates. The total metabolic reaction for *Bombyx mori* larva:




Table 3. Stoichiometry of biosynthesis, catabolism and entire metabolism: v_{pr} is the stoichiometric coefficient of proteins, v_{ip} – the stoichiometric coefficient of lipids, v_{ch} – the stoichiometric coefficient of carbohydrates, v_{bio} – the stoichiometric coefficient of live matter; the stoichiometric coefficients are positive for products and negative for reactants

Process	v_{pr}	v_{ip}	v_{ch}	v_{bio}	$v_{\text{CH}_4\text{N}_2\text{O}}$	$v_{\text{H}_2\text{O}}$	v_{O_2}	v_{CO_2}
Biosynthesis	-1.0000	0.0000	0.0000	1.0000	0.0349	-0.1149	0.0941	-0.0349
Catabolism	0.0000	-0.1600	-0.8400	0	0.0000	0.9937	-1.0674	1.0000
Entire metabolism	-1.0000	-0.0441	-0.2315	1.0000	0.0349	0.1589	-0.2000	0.2407

Table 4 shows thermodynamic properties of biosynthesis, catabolism and entire metabolism of *Bombyx mori* larva. They were calculated with the Hess's law based on the biosynthesis, catabolism and total metabolism reactions, tab. 4, and thermodynamic properties of live matter, tab. 2. These include standard reaction enthalpies, $\Delta_r H^0$, entropies, $\Delta_r S^0$, Gibbs energies, $\Delta_r G^0$, and heat capacities, $\Delta_r C_{p,m}^0$. Standard enthalpy of biosynthesis is 94.06 kJ/C-mol, standard entropy of biosynthesis is 6.59 J/C-molK, standard Gibbs energy of biosynthesis is

92.08 kJ/C-mol and standard heat capacity of biosynthesis is -5.07 J/C-mol K. Standard enthalpy of catabolism is -554.79 kJ/C-mol, standard entropy of catabolism is 22.45 J/C-mol K, standard Gibbs energy of catabolism is -561.41 kJ/C-mol and standard heat capacity of catabolism is 43.15 J/C-mol K. Standard enthalpy of metabolism is -58.80 kJ/C-mol, standard entropy of metabolism is 12.78 J/C-mol K, standard Gibbs energy of metabolism is -62.60 kJ/C-mol and standard heat capacity of metabolism is 6.82 J/C-mol K.

Table 4. Thermodynamic properties of biosynthesis, catabolism and entire metabolism:
 $\Delta_r H^0$ standard reaction enthalpy, $\Delta_r S^0$ standard reaction entropy,
 $\Delta_r G^0$ standard reaction Gibbs energy, $\Delta_r C_{p,m}^0$ standard reaction heat capacity

Name	$\Delta_r H^0$ [kJC-mol ⁻¹]	$\Delta_r S^0$ [JC-mol ⁻¹ K ⁻¹]	$\Delta_r G^0$ [kJC-mol ⁻¹]	$\Delta_r C_{p,m}^0$ [JC-mol ⁻¹ K ⁻¹]
Biosynthesis	94.06	6.59	92.08	-5.07
Catabolism	-554.79	22.45	-561.41	43.15
Entire metabolism	-58.80	12.78	-62.60	6.82

Table 5 gives kinetic properties of biosynthesis, catabolism and metabolism of *Bombyx mori* larva: \dot{q}_{bs} is the biosynthesis heat rate, \dot{q}_{cat} – the catabolism heat rate, r_{cat} – the catabolism reaction rate, and Y – the biomass yield. The biosynthesis heat rate is $2.38 \cdot 10^{-4}$ kW/C-mol, the catabolism heat rate is $-3.87 \cdot 10^{-4}$ kW/C-mol, the catabolism reaction rate is $6.97 \cdot 10^{-7}$ C-mol/C-mols, and the biomass yield is 3.63.

Table 5. Kinetic properties of biosynthesis, catabolism and entire metabolism: \dot{q}_{met} metabolic heat rate, \dot{q}_{bs} – biosynthesis heat rate, \dot{q}_{cat} – catabolism heat rate, r_{bs} – biosynthesis reaction rate, r_{cat} – catabolism reaction rate, Y – biomass yield; the r_{bs} and \dot{q}_{met} values were taken from [111, 112], respectively

\dot{q}_{met} [kW/C-mol ⁻¹]	\dot{q}_{bs} [kW/C-mol ⁻¹]	\dot{q}_{cat} [kW/C-mol ⁻¹]	r_{bs} [kW/C-mol ⁻¹ s ⁻¹]	r_{cat} [kW/C-mol ⁻¹ s ⁻¹]	Y
$-1.49 \cdot 10^{-04}$	$2.38 \cdot 10^{-04}$	$-3.87 \cdot 10^{-04}$	$2.53 \cdot 10^{-06}$	$6.97 \cdot 10^{-07}$	3.63

Table 6 gives reaction affinities and phenomenological coefficients for *Bombyx mori* larva: A_{cat} is the catabolism affinity, A_{bs} – the biosynthesis affinity, L_{CC} – the catabolism-catabolism phenomenological coefficient, L_{BB} – the biosynthesis-biosynthesis phenomenological coefficient, and L_{BC} – the biosynthesis-catabolism (coupling) phenomenological coefficient. They were calculated based on the phenomenological equations [39, 110, 123]. The catabolism affinity is 1.8830 kJ/C-molK, biosynthesis affinity is -0.3088 kJ/C-mol K, catabolism-catabolism phenomenological coefficient is $9.1463 \cdot 10^{-7}$ C-mol K/kJs, biosynthesis-biosynthesis phenomenological coefficient is $1.2048 \cdot 10^{-5}$ C-mol K/kJs and biosynthesis-catabolism phenomenological coefficient is $3.3196 \cdot 10^{-6}$ C-mol K/kJs.

Table 6. Reaction affinities (driving forces) and phenomenological coefficients:
 A_{cat} – the catabolism affinity, A_{bs} – the biosynthesis affinity, L_{CC} – the catabolism-catabolism phenomenological coefficient, L_{BB} – the biosynthesis-biosynthesis phenomenological coefficient, and L_{BC} – the biosynthesis-catabolism (coupling) phenomenological coefficient

A_{cat} [kJC-mol ⁻¹ K ⁻¹]	A_{bs} [kJC-mol ⁻¹ K ⁻¹]	L_{CC} [C-molKk ⁻¹ s ⁻¹]	L_{BB} [C-molKk ⁻¹ s ⁻¹]	L_{BC} [C-molKk ⁻¹ s ⁻¹]
1.8830	-0.3088	$9.1463E \cdot 10^{-07}$	$1.2048 \cdot 10^{-05}$	$3.3196 \cdot 10^{-06}$

Discussion


All organisms originate from the last universal common ancestor (LUCA). Therefore, all organisms interact with their environment and therefore, obey the same laws of chemistry and physics. *Bombyx mori* represents an open thermodynamic system with the property of

growth, which interacts with its surroundings. From the surroundings it takes nutrients and metabolizes them. A part of the substances taken are incorporated into new live matter produced by the organism, which thus represents a growing system. The rest is exported into the environment as CO₂, H₂O, and urea. This is why *Bombyx mori* performs catabolic and anabolic reactions that together make the metabolism.

The empirical formula of *Bombyx mori* is given in tab. 1. The empirical formula of *Bombyx mori* :



The empirical formula of the human organism [101]:



The empirical formula of the *Saccharomyces cerevisiae*, which belongs to eukaryotic microorganisms (fungi) [50]:




The empirical formula of the plant *Zea mays* (corn) [100]:




The empirical formula of the bacterium *Escherichia coli* [124]:




The empirical formula of *Chlorella* microalgae [125]:



The empirical formula of the SARS-CoV-2 virus XBB.1.5 variant nucleocapsid [92]:




The empirical formula of the Ebola virus nucleocapsid [126]:




The differences in elemental composition are a consequence of changes that appeared during evolution, from LUCA to the different species of organisms present today. This paper reports for the first time the empirical formula of an insect. Thus for now it is not possible to make a comparison with other insects.


Knowing empirical formulas enables to define biosynthesis, catabolism and entire metabolism reactions for organisms [39, 40, 51, 52, 127, 128]. The reactions were formulated for the larval stage of *Bombyx mori*, since growth is the most intense in this stage. The biosynthetic (anabolic) reaction needed for growth of *Bombyx mori* larva is



where CH_{1.59}O_{0.32}N_{0.26} is the empirical formula of protein and CH₄N₂O is the formula of urea. In this reaction, proteins are transformed into new live matter, represented by its empirical formula, CH_{1.6803}O_{0.2815}N_{0.1903}. The excess nitrogen is released as urea.


The catabolic reactions for *Bombyx mori* larva start from lipids and carbohydrates as reactants, which are degraded into CO₂ and H₂O:





where $\text{CH}_{1.92}\text{O}_{0.12}$ is the empirical formula of lipids and CH_2O is the empirical formula of carbohydrates. The catabolic processes release energy, which drives the metabolism.

The entire metabolism of *Bombyx mori* is described by the total metabolic reaction. In this reaction the nutrients (proteins, lipids and carbohydrates) are converted into new live matter ($\text{CH}_{1.6803}\text{O}_{0.2815}\text{N}_{0.1903}$), additional product (urea) and catabolic products (CO_2 and H_2O):



The driving force for all chemical reactions is Gibbs energy [39, 110, 123]. The driving force (Gibbs energy) of biosynthesis is 92.08 kJ/C-mol, tab. 4. Gibbs energy of biosynthesis of *Bombyx mori* is positive. This means that the process of biosynthesis on its own is an unfavorable process. In order for biosynthesis to proceed, it must be coupled with a thermodynamically favorable process, to provide the necessary driving force in form of negative Gibbs energy [39, 110, 123]. This process is the catabolism. The driving force for catabolism of *Bombyx mori* is its highly negative Gibbs energy: -554.79 kJ/C-mol, tab. 4. This means that catabolism is highly favorable and provides enough Gibbs energy for both itself and biosynthesis. Therefore, the driving force for the entire metabolism is -62.60 kJ/C-mol. The negative Gibbs energy value means that metabolism is a favorable process and can occur spontaneously in nature.

The driving force for the metabolism of *Bombyx mori* is -62.60 kJ/C-mol. On the other hand, the driving force for metabolism of microorganisms (bacteria) is -500 kJ/C-mol [40, 45, 129]. The driving force of metabolism, $\Delta_{\text{met}}G$, is important since it is proportional to growth rate, r , according to the growth phenomenological equation:

$$r = -\frac{L_g}{T} \Delta_{\text{met}}G \quad (38)$$

(where L_g is the growth phenomenological coefficient) [39, 94, 123]. This means that since the driving force of microorganisms is much greater than that of *Bombyx mori*, microorganisms should have a greater multiplication rate. Indeed, microorganisms can divide every 20 minutes [120]. On the other hand, *Bombyx mori* larvae take days to grow [111].

This research also reports kinetic properties of biosynthesis, catabolism and entire metabolism of *Bombyx mori*, tab. 5. The biosynthesis heat rate is $2.38 \cdot 10^{-4}$ kW/C-mol, while the catabolism heat rate is $-3.87 \cdot 10^{-4}$ kW/C-mol. This means that catabolism can provide enough energy to make entire metabolic process exothermic ($-1.49 \cdot 10^{-4}$ kW/C-mol [112]).

Table 6 gives reaction affinities and phenomenological coefficients for catabolism and biosynthesis of *Bombyx mori*. The phenomenological coefficients allow us to analyze how energy of catabolism is used to make biosynthesis feasible. The rate of catabolism is given by the catabolism phenomenological equation:

$$r_{\text{cat}} = \left(-\frac{L_{\text{CC}}}{T} \Delta_{\text{cat}}G \right) + \left(-\frac{L_{\text{BC}}}{T} \Delta_{\text{bs}}G \right) \quad (39)$$

where L_{CC} is the catabolism-catabolism phenomenological coefficient, L_{BC} the biosynthesis-catabolism phenomenological coefficient, $\Delta_{\text{cat}}G$ Gibbs energy of catabolism and $\Delta_{\text{bs}}G$ Gibbs energy of biosynthesis [39, 123]. Similarly, the rate of biosynthesis is given by the biosynthesis phenomenological equation:

$$r_{\text{bs}} = \left(-\frac{L_{\text{BC}}}{T} \Delta_{\text{cat}} G \right) + \left(-\frac{L_{\text{BB}}}{T} \Delta_{\text{bs}} G \right) \quad (40)$$

where L_{BB} is the biosynthesis-biosynthesis phenomenological coefficient [39, 123]. The biosynthesis reaction of *Bombyx mori* has a positive Gibbs energy change of +92.08 kJ/C-mol. The biosynthesis-biosynthesis phenomenological coefficient is $1.2048 \cdot 10^{-5}$ C-mol K/kJs. Due to the negative sign in the biosynthesis phenomenological equation, this means that the biosynthesis reaction is not a favorable process. However, biosynthesis is coupled to catabolism, which is described through the biosynthesis-catabolism phenomenological coefficient, which is $3.3196 \cdot 10^{-6}$ C-mol K/kJs. Gibbs energy of catabolism is -561.41 kJ/C-mol. This energy hence makes biosynthesis possible. The biosynthesis-catabolism coupling also regulates the rate of catabolism. The catabolism-catabolism phenomenological coefficient is $9.1463 \cdot 10^{-7}$ C-mol K/kJs and Gibbs energy of catabolism is -561.41 kJ/C-mol. This means that the catabolic processes should proceed very fast. However, since Gibbs energy of biosynthesis is positive (+92.08 kJ/C-mol), it decreases the catabolism rate according to the catabolism phenomenological equation. Therefore, catabolism makes biosynthesis feasible and biosynthesis regulates the rate of catabolism.

Conclusions

The empirical formula is reported for the first time for an insect: *Bombyx mori* (domestic silk moth). The empirical formula of *Bombyx mori* is $\text{CH}_{1.6803}\text{O}_{0.2815}\text{N}_{0.1903}\text{S}_{0.005122}$. Biosynthesis, catabolic and total metabolic reactions were formulated for *Bombyx mori*. They show how nutrients that provide energy and elements are transformed into new live matter and other metabolic products.

Driving force (Gibbs energy) of metabolism is reported for the first time for an insect. Gibbs energy of metabolism of *Bombyx mori* is -62.60 kJ/C-mol. The negative Gibbs energy of metabolism means that metabolism and growth of *Bombyx mori* are thermodynamically favorable processes that can occur spontaneously in nature. Furthermore, driving force (Gibbs energy) of metabolism of *Bombyx mori* is much less negative than that of microorganisms (-500 kJ/C-mol). This can explain from the bioenergetic perspective why the growth rate of insects is much lower than that of microorganisms.

Driving forces (Gibbs energies) of biosynthesis and catabolism are reported for the first time for an insect. Gibbs energy of biosynthesis of *Bombyx mori* is +92.08 kJ/C-mol, while Gibbs energy of catabolism is -561.41 kJ/C-mol. Therefore, due to the positive Gibbs energy, biosynthesis is not a favorable process. However, the highly negative Gibbs energy of catabolism provides enough driving force to make biosynthesis feasible, according to the biosynthesis phenomenological equation.

Acknowledgment

This work was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Grant No. 451-03-47/2023-01/200026).

Author statement

Marko E. Popović: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data Curation, Writing – Original Draft, Writing – Review and Editing.

Marijana Pantović Pavlović: Validation, Resources, Writing – Review and Editing, Funding acquisition.

Gavrilo Šekularac: Validation, Resources, Writing – Review and Editing, Funding acquisition.

References

- [1] Carnot, L., Essai sur les machines en général, (in French, Essay On Machines In General), De l'imprimerie de Defay, Dijon, France, 1786, ISBN-13: 978-1147666625
- [2] Carnot, L., Principes fondamentaux de l'équilibre et du mouvement, (in French, Fundamental Principles of Equilibrium and Movement), De l'imprimerie de Crapelet, Paris, France, 1803, ISBN: 2016170190
- [3] Carnot, S., Réflexions sur la puissance motrice du feu et sur les machines propres à développer cette puissance, (in French, Reflections on the Motive Power of Fire and on Machines Fitted to Develop That Power), Bachelier, Paris, France, 1824, ISBN-13: 978-0486446417
- [4] Lavoisier, A. L., Marquis de Laplace, P. S., Mémoire sur la chaleur: Lû à l'Académie royale des sciences, le 28 Juin 1783, (in French, Memoir on Heat Read to the Royal Academy of Sciences, 28 June 1783) De l'Imprimerie royale, Paris, France, 1783
- [5] Lavoisier, A. L., DeLaplace, P. S., Memoir on heat read to the royal academy of sciences, 28 June 1783, *Obesity Research*, 2 (1994), 2, 189-202
- [6] Clausius, R., *The Mechanical Theory of Heat – With its Applications to the Steam Engine and to Physical Properties of Bodies*, John van Voorst, London, UK, 1867
- [7] Clausius, R., On a Mechanical Theorem Applicable to Heat, *Philosophical Magazine Series 4*, 40 (1870), 265, pp. 122-127
- [8] Clausius, R., On Different Forms of the Fundamental Equations of the Mechanical Theory of Heat and Their Convenience for Application, in: *The Second Law of Thermodynamics*, (Ed. J. Kestin), Downen, Hutchinsonson and Ross, Inc., Stroudsburg, Penn., USA, 1976, pp. 162-193
- [9] Muller, I., *A History of Thermodynamics: The Doctrine of Energy and Entropy*, Springer, Berlin, Germany, 2010
- [10] Popovic, M., Living Organisms from Prigogine's Perspective: An Opportunity to Introduce Students to Biological Entropy Balance, *Journal of Biological Education*, 52 (2018), 3, pp. 294-300
- [11] Schrodinger, E., *What is Life? The Physical Aspect of the Living Cell*, Cambridge University Press, Cambridge, UK, 1944
- [12] Prigogine, I., Wiame, J. M., Biologie et thermodynamique des phénomènes irréversibles, (in French, Biology and Thermodynamics of Irreversible Phenomena), *Experientia*, 2 (1946), pp. 451-453
- [13] Prigogine, I., Time, Structure, And Fluctuations, *Science*, 201 (1978), 4358, pp. 777-785
- [14] Prigogine, I., Nobel lecture: Time, Structure and Fluctuations, 1977, <https://www.nobelprize.org/prizes/chemistry/1977/prigogine/lecture/>
- [15] Prigogine, I., Etude thermodynamique des phénomènes irréversibles, (in French, Thermodynamic Study of Irreversible Phenomena), Dunod, Paris, France, 1947, WorldCat ID421502786
- [16] Glansdorff, P., Prigogine, I., *Thermodynamic Theory of Structure, Stability and Fluctuations*, Wiley, Hoboken, N. J., USA, 1971
- [17] Kondepudi, D., Prigogine, I., *Modern Thermodynamics: From Heat Engines to Dissipative Structures*, John Wiley and Sons, Hoboken, N. J., USA, 2014
- [18] Prigogine, I., Rice, S. A., *Proteins: A Theoretical Perspective of Dynamics, Structure, and Thermodynamics*, John Wiley and Sons, Hoboken, N. J., USA, 1991, Vol. 71 (Vol. 86)
- [19] Prigogine, I., Nicolis, G., Biological Order, Structure and Instabilities1, *Quarterly Reviews of Biophysics*, 4 (1971), 2-3, pp. 107-148
- [20] Morowitz, H. J., et al., The Origin of Intermediary Metabolism, *Proceedings of the National Academy of Sciences*, 97 (2000), 14, pp. 7704-7708
- [21] Morowitz, H., The Emergence of Complexity, *Complexity*, 1 (1995), 1, pp. 4-5
- [22] Morowitz, H. J., *Beginnings of Cellular Life: Metabolism Recapitulates Biogenesis*, Yale University Press, New Haven, Ct., USA, 1992
- [23] Morowitz, H. J., et al., The Chemical Logic of a Minimum Protocell, *Origins Life Evol Biosphere*, 18 (1988), Sept., pp. 281-287
- [24] Morowitz, H. J., The High Cost of Being Human, *The New York Times*, (1976) 15, 45
- [25] Morowitz, H. J., *Energy Flow in Biology: Biological Organization as a Problem in Thermal Physics*, Academic Press, New York, N. Y., USA, 1968
- [26] Morowitz, H. J., Some Order-Disorder Considerations in Living Systems, *Bulletin of Mathematical Biophysics*, 17 (1955), June, pp. 81-86
- [27] Hansen, L. D., et al., Transformation of Matter in Living Organisms during Growth and Evolution, *Biophysical Chemistry*, 271 (2021), 106550
- [28] Hansen, L. D., et al., Laws of Evolution Parallel the Laws of Thermodynamics, *The Journal of Chemical Thermodynamics*, 124 (2018), Sept., pp. 141-148

- [29] Hansen, L. D., et al., Biological calorimetry and The Thermodynamics of the Origination and Evolution of Life, *Pure and Applied Chemistry*, 81 (2009), 10, pp. 1843-1855
- [30] Hansen, L. D., et al., From Biochemistry to Physiology: The Calorimetry Connection, *Cell Biochemistry and Biophysics*, 49 (2007), 2, pp. 125-140
- [31] Hansen, L. D., et al., Use of Calorespirometric Ratios, Heat per CO₂ and Heat per O₂, to Quantify Metabolic Paths and Energetics of Growing Cells, *Thermochimica Acta*, 422 (2004), 1-2, pp. 55-61
- [32] Hansen, L. D., et al., Kinetics of Plant Growth and Metabolism, *Thermochimica Acta*, 388 (2002), 1-2, pp. 415-425
- [33] Hansen, L. D., et al., Responses of Plant Growth and Metabolism to Environmental Variables Predicted from Laboratory Measurements, *Proceedings*, USDA Forest Service Proceedings RMRS-P-00, 2001, pp. 1-6
- [34] Ellingson, D., et al., Determination of the Enthalpy Change for Anabolism by Four Methods, *Thermochimica Acta*, 400 (2003), 1-2, pp. 79-85
- [35] Russell, D. J., Hansen, L. D., Calorimeters for Biotechnology, *Thermochimica Acta*, 445 (2006), 2, pp. 151-159
- [36] Macfarlane, C., et al., Application of an Enthalpy Balance Model of The Relation Between Growth and Respiration Temperature Acclimation of Eucalyptus Globulus Seedlings, *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 269 (2002), 1499, pp. 1499-1507
- [37] Smith, B. N., et al., Time, Plant Growth, Respiration, and Temperature, in: *Handbook of Plant and Crop Physiology*, (Ed. M. Pessaraki), CRC Press, Boca Raton, Fla., USA, 2001, pp. 23-34
- [38] Robinson, C. M., et al., Temperature Response of Metabolic Activity of an Antarctic Nematode, *Biology*, 12 (2023), 1, 109
- [39] Von Stockar, U., Live Cells as Open Non-Equilibrium Systems, in: *Biothermodynamics: The Role of Thermodynamics in Biochemical Engineering*, (Ed. U. von Stockar), EPFL Press, Lausanne, Switzerland, 2013, pp. 399-421
- [40] Von Stockar, U., Biothermodynamics of Live Cells: Energy Dissipation and Heat Generation in Cellular Structures. in: *Biothermodynamics: The Role of Thermodynamics in Biochemical Engineering*, (Ed. U. von Stockar), EPFL Press, Lausanne, Switzerland, 2013, pp. 475-534
- [41] Von Stockar, U., et al., Thermodynamic Analysis of Metabolic Pathways, Biothermodynamics of Live Cells: Energy Dissipation and Heat Generation in Cellular Structures. in: *Biothermodynamics: The Role of Thermodynamics in Biochemical Engineering*, (Ed. U. von Stockar), EPFL Press, Lausanne, Switzerland, 2013, pp. 581-604
- [42] Von Stockar, U., et al., Can Microbial Growth Yield Be Estimated Using Simple Thermodynamic Analogies to Technical Processes, *Chemical Engineering and Processing, Process Intensification*, 47 (2008), 6, pp. 980-990
- [43] Von Stockar, U., et al., Thermodynamics of Microbial Growth and Metabolism: An Analysis of the Current Situation, *Journal of Biotechnology*, 121 (2006), 4, pp. 517-533
- [44] Von Stockar, U., et al., Thermodynamic Considerations in Constructing Energy Balances, *Biochimica et Biophysica Acta*, 1183 (1993), 2, pp. 221-240
- [45] Von Stockar, U., Liu, J., Does Microbial Life Always Feed on Negative Entropy, Thermodynamic Analysis of Microbial Growth, *Biochimica et Biophysica Acta*, 1412 (1999), 3, pp. 191-211
- [46] Maskow, T., von Stockar, U., How Reliable Are Thermodynamic Feasibility Statements Of Biochemical Pathways, *Biotechnology and Bioengineering*, 92 (2005), 2, pp. 223-230
- [47] Schill, N. A., et al., Thermodynamic Analysis of Growth of Methanobacterium Thermoautotrophicum, *Biotechnology and Bioengineering*, 64 (1999), 1, pp. 74-81
- [48] Assael, M. J., et al., *Commonly Asked Questions in Thermodynamics*, 2nd ed., CRC Press, Boca Raton, Fla., 2022
- [49] Battley, E. H., A theoretical Study of the Thermodynamics of Microbial Growth Using Saccharomyces Cerevisiae and A Different Free Energy Equation, *The Quarterly Rev. of Bio.*, 88 (2013), 2, pp. 69-96
- [50] Battley, E. H., An Empirical Method for Estimating the Entropy of Formation and the Absolute Entropy of Dried Microbial Biomass for Use in Studies on the Thermodynamics of Microbial Growth, *Thermochimica Acta*, 326 (1999), 1-2, pp. 7-15
- [51] Battley, E. H., The Thermodynamics of Microbial Growth, in: *Handbook of Thermal Analysis and Calorimetry, From Macromolecules to Man* (Ed. E. B. Kemp) Elsevier, Amsterdam, Netherlands, 1999, Vol. 4, pp. 219-235
- [52] Battley, E. H., The Development of Direct and Indirect Methods for the Study of The Thermodynamics of Microbial Growth, *Thermochimica Acta*, 309 (1998), 1-2, pp. 17-37

- [53] Battley, E. H., *et al.*, Heat Capacity Measurements from 10 To 300 K and Derived Thermodynamic Functions of Lyophilized Cells of *Saccharomyces Cerevisiae* Including the Absolute Entropy and the Entropy of Formation at 298.15 K, *Thermochimica Acta*, 298 (1997), 1-2, pp. 37-46
- [54] Battley, E. H., On the Thermodynamics of Autotrophic and Heterotrophic Growth of *Pseudomonas Saccharophila*, *Canadian Journal of Microbiology*, 42 (1996), 1, pp. 38-45
- [55] Battley, E. H., The Advantages and Disadvantages of Direct and Indirect Calorimetry, *Thermochimica Acta*, 250 (1995), 2, pp. 337-352
- [56] Battley, E. H., Calculation of Entropy Change Accompanying Growth of *Escherichia Coli* K-12 on Succinic Acid, *Biotechnology and Bioengineering*, 41 (1993), 4, pp. 422-428
- [57] Battley, E. H., On the Enthalpy of Formation of *Escherichia Coli* K-12 Cells, *Biotechnology and Bioengineering*, 39 (1992), 1, pp. 5-12
- [58] Battley, E. H., Calculation of Thermodynamic Properties of Protein in *Escherichia Coli* K-12 Grown on Succinic Acid, Energy Changes Accompanying Protein Anabolism, and Energetic Role of Atp in Protein Synthesis, *Biotechnology and Bioengineering*, 40 (1992), 2, pp. 280-288
- [59] Battley, E. H., Stone, J. R., A Comparison of Values for The Entropy and the Entropy of Formation of Selected Organic Substances of Biological Importance in the Solid State, As Determined Experimentally Or Calculated Empirically, *Thermochimica Acta*, 349 (2000), 1-2, pp. 153-161
- [60] Maskow, T., Pauffler, S., What Does Calorimetry and Thermodynamics of Living Cells Tell Us, *Methods*, 76 (2015), Apr., pp. 3-10
- [61] Maskow, T., Babel, W., Thermokinetic Description of Anaerobic Growth of *Halomonas Halodenitrificans* Using a Static Microcalorimetric Ampoule Technique, *Journal of Biotechnology*, 101 (2003), 3, pp. 267-274
- [62] Maskow, T., Babel, W., Calorimetric Analysis of Microorganisms in Transient Growth States to Quantify Changes of Metabolic Fluxes in Response to Nutrient Deficiencies and Osmostress, *Thermochimica Acta*, 382 (2002), 1-2, pp. 229-237
- [63] Maskow, T., Babel, W., Calorimetrically Obtained Information about the Efficiency of Ectoine Synthesis From Glucose in *Halomonas Elongata*, *Biochimica et Biophysica Acta (BBA)-General Subjects*, 1527 (2001), 1-2, pp. 4-10
- [64] Maskow, T., Kemp, R.B., Bioprocess Engineering-Bioprocess Analysis through Calorimetry and Biothermodynamics, *Encyclopedia of Life Support Systems, Biotechnology*, IV (2008), pp. 181-191
- [65] Maskow, T., Kleinstaubert, S., Carbon and Energy Fluxes during Haloadaptation of *Halomonas* sp. EF11 Growing on Phenol, *Extremophiles*, 8 (2004), Jan., pp. 133-141
- [66] Maskow, T., Harms, H., Real Time Insights into Bioprocesses Using Calorimetry: State of the Art and Potential, *Engineering in Life Sciences*, 6 (2006), 3, pp. 266-277
- [67] Maskow, T., *et al.*, Rapid Analysis of Bacterial Contamination of Tap Water Using Isothermal Calorimetry, *Thermochimica Acta*, 543 (2012), Sept., pp. 273-280
- [68] Duong, H. L., *et al.*, Applicability and Information Value of Biocalorimetry for The Monitoring of Fungal Solid-State Fermentation of Lignocellulosic Agricultural by-Products, *New Biotechnology*, 66 (2022), Jan., pp. 97-106
- [69] Mariana, F., *et al.*, Isothermal Titration Calorimetry – A New Method for the Quantification of Microbial Degradation of Trace Pollutants, *Journal of Microbiological Methods*, 82 (2010), 1, pp. 42-48
- [70] Hoffmann, P., *et al.*, A Thermodynamic Investigation of the Glucose-6-Phosphate Isomerization, *Biophysical Chemistry*, 195 (2014), Dec., pp. 22-31
- [71] Skene, K., Life's a Gas: A Thermodynamic Theory of Biological Evolution, *Entropy*, 17 (2015), 12, pp. 5522-5548
- [72] Barros, N., Thermodynamics of Soil Microbial Metabolism: Applications and Functions, *Applied Sciences*, 11 (2021), 11, 4962
- [73] Barros, N., *et al.*, Calorimetry and Soil, *Thermochimica Acta*, 458 (2007), 1-2, pp. 11-17
- [74] Barros, N., Feijoo, S., A combined Mass and Energy Balance to Provide Bioindicators of Soil Microbiological Quality, *Biophysical Chemistry*, 104 (2003), 3, pp. 561-572
- [75] Barros Pena, N., Calorimetry and Soil Biodegradation: Experimental Procedures and Thermodynamic Models, in: *Toxicity and Biodegradation Testing*, (ed. Bidoia *et al.*), Springer, New York, USA, 2018, pp. 123-145
- [76] Seppelt, A., *et al.*, Thermodynamic Characterization of LF, H, and Mineral Soil Layers from Oak Forest Ecosystems: Exploring the Role of Proximate Analysis, *Environmental Research*, 222 (2023), 115310
- [77] Lucia, U., *et al.*, Thermodynamics and SARS-CoV-2: Neurological Effects in post-Covid 19 Syndrome, *Atti della Accademia Peloritana dei Pericolanti*, 99 (2021), 2, A3
- [78] Lucia, U., *et al.*, Seebeck-like Effect in SARS-CoV-2 Biothermodynamics, *Atti della Accademia Peloritana dei Pericolanti-Classe di Scienze Fisiche, Matematiche e Naturali*, 98 (2020), 2, 6

- [79] Lucia, U., *et al.*, Entropy-Based Pandemics Forecasting, *Frontiers in Physics*, 8 (2020), 274
- [80] Lucia, U., Bioengineering Thermodynamics: An Engineering Science for Thermodynamics of Biosystems, *International Journal of Thermodynamics*, 18 (2015), 4, pp. 254-265
- [81] Lucia, U., Bioengineering Thermodynamics of Biological Cells Theoretical Biology and Medical Modelling, 12 (2015), 1, pp.1-16
- [82] Lucia, U., Entropy Generation Approach to Cell Systems, *Physica A: Statistical Mechanics and Its Applications*, 406 (2014), July, pp. 1-11
- [83] Grisolia, G., *et al.*, Biomethanation of Rice Straw: A Sustainable Perspective for the Valorisation of a Field Residue in the Energy Sector, *Sustainability*, 14 (2022), 9, 5679
- [84] Kaniadakis, G., *et al.* The κ -Statistics Approach to Epidemiology, *Sci. Rep.*, 10 (2020), 19949
- [85] Ongel, M. E., *et al.*, Thermodynamic Assessment of the Effects of Intermittent Fasting and Fatty Liver Disease Diets on Longevity, *Entropy*, 25 (2023), 2, 227
- [86] Yildiz, C., Ozilgen, M., Species-Specific Biological Energy Storage and Reuse, *Energy Storage*, 4 (2022), 6, e382
- [87] Yildiz, C., Ozilgen, M., Why Brain Functions May Deteriorate with Aging: A Thermodynamic Evaluation, *International Journal of Exergy*, 37 (2022), 1, pp. 87-101
- [88] Semercioz, A. S., Ozilgen, M., How Much Energy Is Needed to Keep the Stored Cellular Mass and Energy Free of Errors: Case Studies, *Energy Storage*, 5 (2023), 1, e389
- [89] Ozilgen, M., Exergetic Growth Efficiency of *Kluyveromyces Fragilis* on Complex Organic and Minimal Inorganic Media, *International Journal of Exergy*, 40 (2023), 3, 336-346
- [90] Yildiz, C., *et al.*, Organisms Live at Far-From-Equilibrium with Their Surroundings While Maintaining Homeostasis, *Importing Exergy and Exporting Entropy*, *Int. J. of Exergy*, 31 (2020), 3, pp. 287-301
- [91] Ozilgen, M., Yilmaz, B., The COVID-19 Disease Causes an Energy Supply Deficit in a Patient, *International Journal of Energy Research*, 45 (2021), 2, 1157
- [92] Popovic, M. E., The XBB.1.5 Kraken Cracked: Gibbs Energies of Binding and Biosynthesis of the XBB.1.5 variant of SARS-CoV-2, *Microbiological Research*, 270 (2023), 127337
- [93] Popovic, M., Never Ending Story, Evolution of SARS-CoV-2 Monitored through Gibbs Energies of Biosynthesis and Antigen-Receptor Binding of Omicron BQ.1, BQ.1.1, XBB and XBB.1 Variants, *Microbial Risk Analysis*, 23 (2023), 100250
- [94] Popovic, M., Biothermodynamics of Viruses from Absolute Zero (1950) to Virothermodynamics (2022), *Vaccines*, 10 (2022), 12, 2112
- [95] Popovic, M., Thermodynamic Properties of Microorganisms: Determination and Analysis of Enthalpy, Entropy, and Gibbs Free Energy of Biomass, Cells and Colonies of 32 Microorganism Species, *Heliyon*, 5 (2019), 6, e01950
- [96] Popovic, M. E., *et al.*, Upcoming Epidemic Storm: Empirical Formulas, Biosynthesis Reactions, Thermodynamic Properties and Driving Forces of Multiplication of the Omicron XBB. 1.9. 1, XBF and XBB. 1.16 (Arcturus) Variants of SARS-CoV-2, *Microbial Risk Analysis*, 25 (2023), 100273
- [97] Popovic, M., *et al.*, Ghosts of the Past: Elemental Composition, Biosynthesis Reactions and Thermodynamic Properties of Zeta P.2, Eta B.1.525, Theta P.3, Kappa B.1.617.1, Iota B.1.526, Lambda C.37 and Mu B.1.621 Variants of SARS-CoV-2, *Microbial Risk Analysis*, 24 (2023), 100263
- [98] Popovic, M., *et al.*, The COVID Infection in 4 Steps: Thermodynamic Considerations Reveal How Viral Mucosal Diffusion, Target Receptor Affinity and Furin Cleavage Act in Concert to Drive the Nature and Degree of Infection in Human COVID-19 Disease, *Heliyon*, 9 (2023), 6, e17174
- [99] Popovic, M., *et al.*, Elemental Composition, Heat Capacity from 2-300 K and Derived Thermodynamic Functions of 5 Microorganism Species, *Journal of Biotechnology*, 331 (2021), Apr., pp. 99-107
- [100] Popovic, M., Minceva, M., Standard Thermodynamic Properties, Biosynthesis Rates, and the Driving Force of Growth of Five Agricultural Plants, *Frontiers in Plant Science*, 12 (2021), 671868
- [101] Popovic, M. E., Minceva, M., Thermodynamic Properties of Human Tissues, *Thermal Science*, 24 (2020), 6B, pp. 4115-4133
- [102] Popovic, M., The SARS-CoV-2 Strain Wars Continues: Chemical And Thermodynamic Characterization of Live Matter and Biosynthesis of Omicron BN.1, CH.1.1 and XBC Variants, *Microbial Risk Analysis*, 24 (2023), 100260
- [103] Popovic, M., The SARS-CoV-2 Hydra, a Monster from the 21st Century: Thermodynamics of the BA.5.2 and BF.7 Variants, *Microbial Risk Analysis*, 23 (2023), 100249
- [104] Popovic, M., Thermodynamics of Bacteria-Phage Interactions: T4 and Lambda Bacteriophages, and *E. Coli* Can Coexist in Natural Ecosystems Due to the Ratio of Their Gibbs Energies of Biosynthesis, *Thermal Science*, 27 (2023), 1, pp. 411-431

- [105] Degueldre, C., Single Virus Inductively Coupled Plasma Mass Spectroscopy Analysis: A Comprehensive Study, *Talanta*, 228 (2021), 122211
- [106] Simsek, B., et al., How Much Energy Is Stored in SARS-CoV-2 and Its Structural Elements, *Energy Storage*, 4 (2021), 2, e298
- [107] Yilmaz, B., et al., Energetic and exergetic Costs of COVID-19 Infection on the Body of a Patient, *International Journal of Exergy*, 32 (2020), 3, pp. 314-327
- [108] Srivastava, S., et al., Nutritional Quality of Leaves of Some Genotypes of Mulberry (*Morus alba*), *International Journal of Food Sciences and Nutrition*, 57 (2006), 5-6, pp. 305-313
- [109] Wang, Z. M., et al., Five-Level Model: Reconstruction of Body Weight at Atomic, Molecular, Cellular, and Tissue-System Levels from Neutron Activation Analysis, in: *Human Body Composition*, Basic Life Sciences, (Eds: K.J. Ellis and J. D. Eastman), Springer, Boston, Mass., USA, 1993, Vol. 60, pp. 125-128
- [110] Balmer, R.T., *Modern Engineering Thermodynamics*, Academic Press, Cambridge, Mass., 2010
- [111] Konala, N., et al., The Effect of Bovine Milk on The Growth of *Bombyx mori*, *Journal of Insect Science*, 13 (2013), 1, 98
- [112] Makarieva, A. M., et al., Mean Mass-Specific Metabolic Rates Are Strikingly Similar Across Life's Major Domains: Evidence for Life's Metabolic Optimum, *Proceedings of the National Academy of Sciences of the United States of America*, 105 (2008), 44, pp. 16994-16999
- [113] Popovic, M., Atom Counting Method for Determining Elemental Composition of Viruses and Its Applications in Biothermodynamics and Environmental Science, *Computational Biology and Chemistry*, 96 (2022), 107621
- [114] Patel, S. A., Erickson, L. E., Estimation of Heats of Combustion of Biomass from Elemental Analysis Using Available Electron Concepts, *Biotechnology and Bioengineering*, 23 (1981), 9, pp. 2051-2067
- [115] Hurst, J. E., Harrison, B. K., Estimation of Liquid and Solid Heat Capacities Using a Modified Kopp's Rule, *Chemical Engineering Communications*, 112 (1992), 1, pp. 21-30
- [116] Ozilgen, M., Sorguven, E., *Biothermodynamics: Principles and Applications*, CRC Press, Boca Raton, Fla., USA, 2017
- [117] Atkins, P. W., de Paula, J., *Physical Chemistry for the Life Sciences*, 2nd ed., W. H. Freeman and Company, London, UK, 2011
- [118] Atkins, P. W., de Paula, J., *Physical Chemistry: Thermodynamics, Structure, and Change*, 10th ed., W. H. Freeman and Company, London, UK, 2014
- [119] Berg, J. M., et al., *Biochemistry*, 5th ed., Freeman, New York, USA, 2002
- [120] Riedel, S., et al., *Jawetz, Melnick and Adelbergs Medical Microbiology*, 28th ed., McGraw-Hill Education, New York, USA, 2019
- [121] Privalov, P. L., *Microcalorimetry of Macromolecules: The Physical Basis of Biological Structures*, John Wiley and Sons, Hoboken, N. J., 2012
- [122] Sarge, S. M., et al., *Calorimetry: Fundamentals, Instrumentation and Applications*, John Wiley and Sons, Hoboken, N. J., 2014.
- [123] Demirel, Y., *Non-Equilibrium Thermodynamics: Transport and Rate Processes in Physical, Chemical and Biological Systems*, 3rd ed., Elsevier, Amsterdam, The Netherlands, 2014
- [124] Duboc, P., et al., Quantitative Calorimetry and Biochemical Engineering, in: *Handbook of thermal analysis and calorimetry: From Macromolecules to Man* (Ed. R.B. Kemp), Elsevier, Amsterdam, The Netherlands, 1999, pp. 267-365
- [125] Manahan, S., Manahan, S. E., *Environmental Chemistry*, 9th ed., CRC Press, Boca Raton, Fla., USA, 2009
- [126] Popovic, M., Why Does not Ebola Virus Cause Pandemics Like SARS-CoV-2, *Microbial Risk Analysis*, 22 (2022), 100236
- [127] Popovic, M., Beyond COVID-19: Do Biothermodynamic Properties allow Predicting the Future Evolution of SARS-CoV-2 Variants, *Microbial Risk Analysis*, 22 (2022), 100232
- [128] Popovic, M., Strain Wars 3: Differences in Infectivity and Pathogenicity between Delta and Omicron Strains of SARS-CoV-2 Can Be Explained by Thermodynamic and Kinetic Parameters of Binding and Growth, *Microbial Risk Analysis*, 22 (2022), 100217
- [129] Liu, J. S., et al., A Comparison of Various Gibbs Energy Dissipation Correlations for Predicting Microbial Growth Yields, *Thermochimica Acta*, 458 (2007), 1-2, pp. 38-46