

NUMERICAL SIMULATION STUDY ON THE ACTION LAW OF LOW-TEMPERATURE DRILLING FLUID INTRUSION INTO ROCK FORMATION IN POLAR DRILLING

by

**Jintang WANG^{a,b,*}, Lei LIU^{a,b}, Ke ZHAO^{a,b}, Tong GAO^{a,b}, Jingping LIU^{a,b}, Kaihe LV^{a,b},
Bo LIAO^{a,b}, Jinsheng SUN^{a,b}**

^a State Key Laboratory of Unconventional Oil & Gas Development, China University of Petroleum (East China), Ministry of Education, Qingdao 266580, China

^b School of Petroleum Engineering, China University of Petroleum (East China), Qingdao 266580, China

In this paper, based on the experimental parameters of polar permafrost formation, the mathematical module, solid mechanics module, and Darcy's law module of COMSOL Multiphysics software are used to realize the coupled solution of partial differential equations of temperature field, moisture field, and stress field considering the ice-water phase transition and moisture convection, and the relationship between the permafrost moisture convection and temperature change is analyzed by using the heat-flow-solid-moisture mutual coupling, which reveals that The law of action of low-temperature intrusion into rock formation by polar drilling.

Key words: Polar permafrost, Cryogenic drilling mud invasion, Wellbore–reservoir coupling, open problems, numerical simulation

Introduction

At this stage, with the continuous improvement of drilling technology, researchers are gradually focusing on unconventional reservoir oil and gas exploration. Since the United States explored rich oil and gas resources in the Arctic region in 2008, various countries have also carried out oil and gas exploration and development of the Arctic Antarctic and other polar tundra[1,2]. However, there are few reports of successful drilling cases in the polar regions, as the drilling problems related to the polar tundra are still being explored[3].

Most of the polar drilling is permafrost rock samples[4]. Permafrost is a temperature-sensitive porous elastic-plastic material; when the low-temperature drilling fluid invades the contrasting permafrost layer due to the pressure and temperature changes, it will make the formation of internal phase changes. Since permafrost has the property of freezing, swelling, thawing, and sinking, the phase change of water will lead to the redistribution of stress and strain in the formation, thus affecting the stability performance of the well wall[5,6].

Aiming at the above problems, this paper carries out the secondary development by mathematically analyzing the temperature field, pressure field, moisture field, and strain field of the

* Corresponding author; e-mail: wangjintang@upc.edu.cn

polar permafrost formation in the process of drilling fluid intrusion, using the coefficient-type partial differential equations module (PDE) in COMSOL Multiphysics finite element software[7,8]. According to the thermodynamic properties of permafrost, the changes in the temperature field, pressure field, and stress field of the reservoir by the temperature intrusion of low-temperature drilling fluid into the polar formation are analyzed, and a coupled model of heat-fluid-solid multiphysics field is established[9,10]. The interaction law between the low-temperature drilling fluid and the contrasting permafrost rock layer is revealed using simulation numerical modeling, which provides a relevant theoretical basis for the subsequent drilling exploration and exploitation of the polar permafrost layer[11].

Multi-physics field modeling

Using a set of multi-physics control equations for temperature, moisture, pressure, and stress fields, a model was built using COMSOL Multiphysics, and the results of numerical simulation calculations were used to reveal the pattern of the drilling fluid when it intruded into the polar rock formation.

Geometric model

The polar permafrost was constructed as a two-dimensional model with a length of 0.2 m and a height of 0.1 m. A user-defined rectangular grid is created using COMSOL. The finite element model is divided into 5000 domain cells and 300 boundary cells, and the computation period is three days with a step size of 1h.

Parameter selection and determination of boundary conditions

The values of thermodynamic parameters of each material of this calculation refer to the contents of references such as "Physics of Permafrost," and the hydraulics parameters are quoted from "Unsaturated Soil Mechanics," etc.

In the model boundary definition, since this model is the drilling fluid intruding into the formation from the left side to interact with the rock, the upper and lower boundaries of the model are regarded as adiabatic and no flux boundaries; there is no heat and mass exchange between the upper and lower sides and the outside world, and the left boundary is the intrusion temperature of the drilling fluid and the boundary load pressure. The material parameters and boundary constraints are added, the right boundary is set as a fixed constraint, and the upper and lower boundaries are set as roll supports. The grid is divided for transient calculations.

Analysis of results

Based on the pre-assumed conditions, this problem can be viewed as a radial displacement problem of drilling fluid in two dimensions, and the results mainly contain temperature field analysis, pressure field analysis, and permafrost strain analysis.

Temperature field analysis

When 0-degree drilling fluid intrudes into the -5-degree formation, the convection and diffusion of heat will cause the temperature within the formation to increase, and the specific changes are shown in Figs. 1 and 2.

As shown in Fig. 1, the temperature of the drilling fluid is higher than that of the permafrost, so the formation temperature gradually increases with time, but the overall temperature change is not drastic because the melting of the permafrost ice layer will absorb the corresponding heat and lead to a further reduction of the formation temperature, and the formation temperature change is minimal after 30 h. The temperature change within 0.2 m near the wellbore is minimal, indicating that under the initial

pump-in pressure of 11 MPa, the formation temperature is 9 MPa. The temperature change within 0.2m near the wellbore is minimal. The change is mainly concentrated in 0.1m outside the wellbore, which indicates that under the initial pumping pressure of 11MPa, the drilling fluid intrusion into the 9MPa formation is not very deep. It can be seen in Fig. 2 that the rate of change of the formation temperature and the final value will be reduced with the increase of the depth of the intrusion of drilling fluid. The temperature at 0.03m increased by 3.48 Within 72h, the temperature at 0.03m increased by 3.48°C. In comparison, the formation temperatures at 0.12m and 0.18m only increased by 1.65°C and 0.41°C, with an average temperature change gradient of 10.8°C/m.




Fig. 1. Temperature cloud of 0°C drilling fluid intrusion into the formation




Fig. 2. Stratigraphic temperature variations at different locations of 0°C drilling fluid intrusion into the formation

Pressure field analysis

When the drilling fluid intrudes into the polar permafrost at 11 MPa, the pressure-driven motion of the drilling fluid approximately follows Darcy's law, which in turn causes the formation pressure to redistribute as well.

As can be seen from the pressure change cloud diagram in Fig. 3 and the pressure change diagram in Fig. 4 of the formation at different locations, the pressure at each formation area gradually

increases with time due to the intrusion of drilling fluid from high-pressure to low-pressure. In the early stage of the change, due to the movement of the water field within the formation from static, the pressure at each location initially fluctuates slightly.




Fig. 3. Pressure cloud of 0°C drilling fluid intrusion into the formation




Fig. 4. The pressure variations at different locations of 0°C drilling fluid intrusion into the formation

Still, in general, it shows an upward trend. As the distance from the wellbore increases, the pressure value and the range of pressure change gradually decrease, and the pressure change within 0.2m near the wellbore is small. The change is mainly concentrated in 0.1m outside the wellbore, which is also coupled with the temperature change. Within 72h, the pressure at 0.03m increased by 1.70 MPa, while the formation temperatures at 0.12m and 0.18m only increased by 0.80 MPa and 0.20 MPa, and the average pressure drop gradient was 4.75 MPa/m. The pressure drop gradient at 0.12m and 0.18m was 0.80 MPa and 0.20 MPa, respectively.

Strain field analysis

When the drilling fluid intrudes into the polar permafrost, it causes a relative increase in the absolute value of the volumetric strain inside the formation due to the permafrost ice-water phase change, as shown in Figs. 5 and 6.




Fig. 5. Volumetric strain cloud of 0°C drilling fluid intrusion into the formation




Fig. 6. Variation of formation volumetric strain at different locations of 0°C drilling fluid intrusion into the formation

From the variation of volumetric strain in Fig. 5 and the variation of volumetric strain in the formation at different locations in Fig. 6, it can be seen that, due to the principle of frost heave and thaw settlement inside permafrost, in permafrost drilling operation, the temperature difference between the drilling fluid and the formation generates thermal stresses. The heat transfer between the two will change the mechanical parameters of the formation, which in turn affects the stability of the formation. When 0°C drilling fluid intrudes into the formation, the ice layer's melting causes the formation volume to decrease, and the volumetric strain is negative. With the influence of phase change and water displacement, the rate of change of the volumetric strain and the final value gradually decreases with the increase of the distance from the wellbore, which indicates that the intrusion range of drilling fluid is still the most drastic in the range of less than 0.1m. It corresponds to the temperature and pressure field mentioned above. Within 72 h, the volumetric strain value at 0.03 m changed by 8.91×10^{-3} , while the formation temperatures at 0.12 m and 0.18 m only increased by 5.88×10^{-3} and 1.15×10^{-3} , and the average volumetric strain gradient was $3.08 \times 10^{-2}/\text{m}$, which was a slight change.

Conclusion

Through COMSOL Multiphysics numerical simulation of low-temperature drilling fluid

intrusion into the polar permafrost, it is concluded that the temperature and pressure of drilling fluid will affect the original state of the formation when the drilling fluid at 0°C with 11MPa intrudes into the polar formation with the actual temperature and pressure of -5°C and 9MPa, the temperature and pressure of the formation will increase to different degrees. With the increased distance from the borehole, the final value and the rate of change will be relatively weakened.

Acknowledgments

This work was supported by the China University of Petroleum (East China) Independent Innovation Research Program Project (27RA2302016), the National Natural Science Foundation of China (No. 52274025 and 51991361), the National Key Research and Development Program of China (2021YFC2800803), the CNPC's Major Science and Technology Projects (No. ZD2019-184-003), and the Key R & D Program of Shandong Province (No. 2020ZLYS07).

References

- [1] Tarnocai, C., *et al.*, Soil Organic Carbon Pools in the Northern Circumpolar Permafrost Region, *Global Biogeochem. Cycles*, 23 (2009), 6, Article ID GB2023
- [2] Vernikovskiy, V.A., *et al.*, Concerning Tectonics and the Tectonic Evolution of the Arctic, *Russian Geology and Geophysics*, 54 (2013), 8, pp.838-858
- [3] Li, Y., *et al.*, Influence of Drilling Fluid Temperature on Borehole Shrinkage During Drilling Operation in Cold Regions, *Journal of Petroleum Science and Engineering*, 190 (2020), 6, Article ID107050
- [4] Eppelbaum, L.V., *et al.*, Well Drilling in Permafrost Regions: Dynamics of the Thawed Zone, *Polar Research*, 38 (2019), 6, Article ID 3351
- [5] Wang, R., *et al.*, Microscopic Molecular Insights into Methane Hydrate Growth on the Surfaces of Clay Minerals: Experiments and Molecular Dynamics Simulations, *Chemical Engineering Journal*, 451 (2023), 1, Article ID 138757
- [6] Zhang, Y., *et al.*, Experimental Study on Ultra-Low Temperature Drilling Fluid in Arctic Permafrost, *Polar Science*, 28 (2021), 6, Article ID 100645
- [7] Wang, Z., *et al.*, Analysis on Wellhead Stability During Drilling Operation in Arctic Permafrost Region, *Proceedings of the ASME 2017 36th International Conference on Ocean Offshore and Arctic Engineering, OMAE2017*, 2017, Article ID 61868
- [8] Deng, X., *et al.*, Application of the Darcy-Stefan Model to Investigate the Thawing Subsidence Around the Wellbore in the Permafrost Region, *Applied Thermal Engineering*, 156 (2019), 6, pp.392-401
- [9] Liao, B., *et al.*, Microscopic Molecular and Experimental Insights into Multi-Stage Inhibition Mechanisms of Alkylated Hydrate Inhibitor, *Energy*, 279 (2023), 9, Article ID 128045
- [10] Wang, X., *et al.*, Coupled Thermal Model of Wellbore and Permafrost in Arctic Regions, *Applied Thermal Engineering*, 123 (2017), 8, pp.1291-1299
- [11] Liao, B., *et al.*, Microscopic Insights into Synergism Effect of Different Hydrate Inhibitors on Methane Hydrate Formation: Experiments and Molecular Dynamics Simulations, *Fuel*, 340 (2023), 5, Article ID 127488

Paper submitted: July 18, 2023

Paper revised: August 21, 2023

Paper accepted: November 18, 2023